

AD-A131 523

LIKELIHOOD RATIO TESTS ON COVARIANCE MATRICES AND MEAN  
VECTORS OF COMPLEX, (U) PITTSBURGH UNIV PA CENTER FOR  
MULTIVARIATE ANALYSIS P R KRISHNAIAN ET AL. MAR 83


1/1

UNCLASSIFIED

AFOSR-TR-83-0692 F49629-82-K-0001

F/G 12/1

NL




END

DATE  
FILED

13 - H A

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS - 1963 - A

AFOSR-TR- 83-0692


(2)

AD A 131 523

**Center for Multivariate Analysis**

**University of Pittsburgh**

Master copy



Approved for public release;  
distribution unlimited.

DTIC  
FEB 27 1983  
E

83 08 19 056

## UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 83-0692</b>	2. GOVT ACCESSION NO. <b>A131523</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "LIKELIHOOD RATIO TESTS ON COVARIANCE MATRICES AND MEAN VECTORS OF COMPLEX MULTIVARIATE NORMAL POPULATIONS AND THEIR APPLICATIONS IN TIME SERIES"	5. TYPE OF REPORT & PERIOD COVERED Technical	
7. AUTHOR(s) J.C. Lee T.C. Chang	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Pittsburgh Pittsburgh, PA 15260	8. CONTRACT OR GRANT NUMBER(s) F49629-82-K-0001	
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NM Bldg. 410 Bolling AFB DC 20332	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5	
14. MONITORING AGENCY NAME & ADDRESS(if different from Controlling Office)	12. REPORT DATE March 1983	
	13. NUMBER OF PAGES 67	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Complex multivariate normal, homogeneity of populations, likelihood ratio tests, multiple independence, multiple time series in the frequency domain, and tables.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper, the authors reviewed the literature on computational aspects of the distributions of the likelihood ratio statistics for testing various hypotheses on the covariance matrices and mean vectors of complex multivariate normal populations. Applications of some of these test procedures in the area of inference on multiple time series in the frequency domain are also discussed. In the Appendix, the authors give tables which are useful in implementation of various likelihood ratio test statistics discussed in this paper.		

DD FORM 1 JAN 73 1473

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LIKELIHOOD RATIO TESTS ON COVARIANCE  
MATRICES AND MEAN VECTORS OF  
COMPLEX MULTIVARIATE NORMAL POPULATIONS  
AND THEIR APPLICATIONS IN TIME SERIES

P. R. Krishnaiah\*

University of Pittsburgh

J. C. Lee

Bell Telephone Laboratories

T. C. Chang

University of Cincinnati

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH  
NO. 83-03  
THE UNIVERSITY OF PITTSBURGH  
APPROVED FOR PUBLIC RELEASE  
W/STC 100-1000  
MATERIALS MANAGEMENT INFORMATION SYSTEM  
ELECTRONIC RECORDING INFORMATION SYSTEM

March 1983

Technical Report No. 83-03

Center for Multivariate Analysis  
University of Pittsburgh  
Ninth Floor, Schenley Hall  
Pittsburgh, PA 15260

\*The work of this author is sponsored by the Air Force Office of Scientific Research under Contract F49629-82-K-0001. Reproduction in whole or in part is permitted for any purpose of the United States Government.

## ABSTRACT

In this paper, the authors reviewed the literature on computational aspects of the distributions of the likelihood ratio statistics for testing various hypotheses on the covariance matrices and mean vectors of complex multivariate normal populations. Applications of some of these test procedures in the area of inference on multiple time series in the frequency domain are also discussed. In the Appendix, the authors give tables which are useful in implementation of various likelihood ratio test statistics discussed in this paper.

### Key words and phrases:

Complex multivariate normal, homogeneity of populations, likelihood ratio tests, multiple independence, multiple time series in the frequency domain, and tables.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DOCUMENT	
UNCLASSIFIED	
INITIAL RELEASE	
PERMIT	
DATE	
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	
61	
62	
63	
64	
65	
66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	
77	
78	
79	
80	
81	
82	
83	
84	
85	
86	
87	
88	
89	
90	
91	
92	
93	
94	
95	
96	
97	
98	
99	
100	
101	
102	
103	
104	
105	
106	
107	
108	
109	
110	
111	
112	
113	
114	
115	
116	
117	
118	
119	
120	
121	
122	
123	
124	
125	
126	
127	
128	
129	
130	
131	
132	
133	
134	
135	
136	
137	
138	
139	
140	
141	
142	
143	
144	
145	
146	
147	
148	
149	
150	
151	
152	
153	
154	
155	
156	
157	
158	
159	
160	
161	
162	
163	
164	
165	
166	
167	
168	
169	
170	
171	
172	
173	
174	
175	
176	
177	
178	
179	
180	
181	
182	
183	
184	
185	
186	
187	
188	
189	
190	
191	
192	
193	
194	
195	
196	
197	
198	
199	
200	
201	
202	
203	
204	
205	
206	
207	
208	
209	
210	
211	
212	
213	
214	
215	
216	
217	
218	
219	
220	
221	
222	
223	
224	
225	
226	
227	
228	
229	
230	
231	
232	
233	
234	
235	
236	
237	
238	
239	
240	
241	
242	
243	
244	
245	
246	
247	
248	
249	
250	
251	
252	
253	
254	
255	
256	
257	
258	
259	
260	
261	
262	
263	
264	
265	
266	
267	
268	
269	
270	
271	
272	
273	
274	
275	
276	
277	
278	
279	
280	
281	
282	
283	
284	
285	
286	
287	
288	
289	
290	
291	
292	
293	
294	
295	
296	
297	
298	
299	
300	
301	
302	
303	
304	
305	
306	
307	
308	
309	
310	
311	
312	
313	
314	
315	
316	
317	
318	
319	
320	
321	
322	
323	
324	
325	
326	
327	
328	
329	
330	
331	
332	
333	
334	
335	
336	
337	
338	
339	
340	
341	
342	
343	
344	
345	
346	
347	
348	
349	
350	
351	
352	
353	
354	
355	
356	
357	
358	
359	
360	
361	
362	
363	
364	
365	
366	
367	
368	
369	
370	
371	
372	
373	
374	
375	
376	
377	
378	
379	
380	
381	
382	
383	
384	
385	
386	
387	
388	
389	
390	
391	
392	
393	
394	
395	
396	
397	
398	
399	
400	
401	
402	
403	
404	
405	
406	
407	
408	
409	
410	
411	
412	
413	
414	
415	
416	
417	
418	
419	
420	
421	
422	
423	
424	
425	
426	
427	
428	
429	
430	
431	
432	
433	
434	
435	
436	
437	
438	
439	
440	
441	
442	
443	
444	
445	
446	
447	
448	
449	
450	
451	
452	
453	
454	
455	
456	
457	
458	
459	
460	
461	
462	
463	
464	
465	
466	
467	
468	
469	
470	
471	
472	
473	
474	
475	
476	
477	
478	
479	
480	
481	
482	
483	
484	
485	
486	
487	
488	
489	
490	
491	
492	
493	
494	
495	
496	
497	
498	
499	
500	
501	
502	
503	
504	
505	
506	
507	
508	
509	
510	
511	
512	
513	
514	
515	
516	
517	
518	
519	
520	
521	
522	
523	
524	
525	
526	
527	
528	
529	
530	
531	
532	
533	
534	
535	
536	
537	
538	
539	
540	
541	
542	
543	
544	
545	
546	
547	
548	
549	
550	
551	
552	
553	
554	
555	
556	
557	
558	
559	
560	
561	
562	
563	
564	
565	
566	
567	
568	
569	
570	
571	
572	
573	
574	
575	
576	
577	
578	
579	
580	
581	
582	
583	
584	
585	
586	
587	
588	
589	
590	
591	
592	
593	
594	
595	
596	
597	
598	
599	
600	
601	
602	
603	
604	
605	
606	
607	
608	
609	
610	
611	
612	
613	
614	
615	
616	
617	
618	
619	
620	
621	
622	
623	
624	
625	
626	
627	
628	
629	
630	
631	
632	
633	
634	
635	
636	
637	
638	
639	
640	
641	
642	
643	
644	
645	
646	
647	
648	
649	
650	
651	
652	
653	
654	
655	
656	
657	
658	
659	
660	
661	
662	
663	
664	
665	
666	
667	
668	
669	
670	
671	
672	
673	
674	
675	
676	
677	
678	
679	
680	
681	
682	
683	
684	
685	
686	
687	
688	
689</	

## 1. Introduction

It is known that certain estimates of spectral density matrices of the stationary and Gaussian multiple time series are distributed approximately as complex Wishart matrices. So, complex multivariate distributions are useful (e.g., see Brillinger (1974), and Hannan (1970)) in the area of inference on multiple time series. These distributions are useful in the area of nuclear physics (see Carmeli (1974)) also.

Wooding (1956) introduced the complex multivariate normal distribution. A complex random vector is said to be distributed as a complex multivariate normal if its real and imaginary parts are distributed jointly as a multivariate normal with a structured covariance matrix. Motivated by applications in time series, Goodman (1963a, b) made a systematic study of the complex multivariate normal distribution and complex Wishart matrix. Since then, James (1964), Khatri (1965), Krishnaiah (1976) and other workers in the field have investigated various aspects of complex multivariate distributions. For a review of the literature on complex multivariate distributions, the reader is referred to Krishnaiah (1976). In this paper, we review the literature on the likelihood ratio tests on mean vectors and covariance matrices of the complex multivariate normal populations as well as some of their applications in the area of inference on multiple time series in the frequency domain.

In Section 2 of this paper, we discuss the complex multivariate normal and complex Wishart matrix. The distribution of the determinant of the complex multivariate beta matrix is discussed in Section 3, whereas Section 4 is devoted to the likelihood ratio test procedure for testing the hypothesis of multiple independence of several sets of variables when their joint distribution is complex multivariate normal. Likelihood ratio tests for the hypothesis of

sphericity and the hypothesis specifying the covariance matrix are discussed in Sections 5 and 6 respectively. In Section 7 we discuss the likelihood ratio test for the homogeneity of the covariance matrices whereas the likelihood ratio test procedure for the homogeneity of several complex multivariate normal populations is discussed in Section 8. Likelihood ratio test procedure specifying the covariance matrix and mean vector is discussed in Section 9. Applications of some test procedures on the covariance matrices of the complex multivariate normal populations to the area of inference on multiple time series in the frequency domain are discussed in Section 10. Various tables useful in implementation of certain likelihood ratio test procedures are given in the Appendix. These tables are constructed by approximating a suitable power of the likelihood ratio statistics with Pearson's Type I distribution by using the first four moments. The accuracy of these tables is found to be good.

## 2. Complex Multivariate Normal and Complex Wishart Distributions

Let  $\underline{z} = \underline{x} + i\underline{y}$  where  $\underline{x}$  and  $\underline{y}$  are of order  $p \times 1$  and  $(\underline{x}', \underline{y}')$  is distributed as  $2p$ -variate normal with mean vector  $(\underline{\mu}_1', \underline{\mu}_2')$  and covariance matrix

$$\underline{C} = \begin{pmatrix} \Sigma_1 & -\Sigma_2 \\ \Sigma_2 & \Sigma_1 \end{pmatrix} \quad (2.1)$$

where  $A'$  denotes the transpose of  $A$ . Then, the distribution of  $\underline{z}$  is known to be the complex multivariate normal distribution with mean vector  $\underline{\mu}$  and covariance matrix  $\Sigma$  where  $\underline{\mu} = \underline{\mu}_1 + i\underline{\mu}_2$  and  $\Sigma = 2(\Sigma_1 - i\Sigma_2)$ . The probability density function (p.d.f.) of  $\underline{z}$  is given by

$$f(\underline{z}) = \frac{1}{\pi^p |\Sigma|} \exp\{-(\underline{z}-\underline{\mu})' \Sigma^{-1} (\underline{z}-\underline{\mu})\} \quad (2.2)$$

whereas the characteristic function of  $\underline{z}$  is

$$\phi(t) = \exp\{it'\underline{\mu} - \frac{1}{4} \bar{t}' \Sigma^{-1} t\} \quad (2.3)$$

where  $t = t_1 + it_2$  and  $\bar{t}$  is the complex conjugate of  $t$ . Wooding (1956) derived expressions for the p.d.f. and characteristic function of  $\underline{z}$ . The

Maximum likelihood estimates of  $\underline{\mu}$  and  $\Sigma$  based on a random sample  $(\underline{z}_1, \dots, \underline{z}_n)$  are known to be

$$\begin{aligned} \hat{\underline{\mu}} &= N^{-1} \sum_{j=1}^N \underline{z}_j \\ \hat{\Sigma} &= N^{-1} \sum_{j=1}^N (\underline{z}_j - \hat{\underline{\mu}})' (\underline{z}_j - \hat{\underline{\mu}}) \end{aligned} \quad (2.4)$$

Also,  $\hat{\mu}$  and  $\hat{\Sigma}$  are distributed independent of each other.

Next, let  $S = N\hat{\Sigma}$ . Then, the distribution of  $S$  is known to be a central complex Wishart matrix with  $n=N-1$  degrees of freedom. The probability density of  $S$  is known (Goodman (1963b)) to be

$$f(S) = \frac{|S|^{n-p} \text{etr}\{-\Sigma^{-1}S\}}{\pi^{p(p-1)/2} \prod_{j=1}^p \Gamma(n-j+1) |\Sigma|^n} \quad (2.5)$$

where  $\text{etr } B$  denotes the exponential of the trace of the matrix  $B$ .

### 3. Distribution of the Determinant of the Complex Multivariate Beta Matrix

In this section, we discuss the distribution of the determinant of the complex multivariate beta matrix. This distribution is useful for testing the hypothesis of the equality of several mean vectors and the equality of two covariance matrices when the underlying distributions are complex multivariate normal. It is also useful in testing the hypothesis  $H_1: \Sigma_{12} = 0$  where  $\Sigma_{12}$  is the covariance between two sets of variables whose joint distribution is complex multivariate normal.

Let  $A_1: p \times p$  and  $A_2: p \times p$  be independently distributed as the central complex Wishart matrices with  $n$  and  $q$  degrees of freedom, and let  $E(A_1/n) = E(A_2/q) = \Sigma$ . Then  $A_1(A_1 + A_2)^{-1}$  is known to be a (central) complex multivariate beta matrix. Now, let

$$U = |A_1(A_1 + A_2)^{-1}|. \quad (3.1)$$

The  $h$ -th moment of  $U$  is known to be

$$E(U^h) = \prod_{j=1}^p \left| \frac{\Gamma(n+h-j+1)}{\Gamma(n-j+1)} \frac{\Gamma(n+q-j+1)}{\Gamma(n+h+q-j+1)} \right|. \quad (3.2)$$

Using the first four moments of  $U^{1/b}$ , Lee, Krishnaiah and Chang (1975) have approximated the distribution of  $U^{1/b}$  with the Pearson's Type I distribution where  $b$  is a properly chosen integer. The constant  $b$  is chosen to be equal to 1 or 2 according as  $M > 20$  or  $M < 20$  where  $M = n - p + 1$ . By making use of this approximation, values of  $c_1$  are computed, where

$$P[C_1 \leq c_1] = (1-\alpha), \quad (3.3)$$

$c_1 = -(2n+q-p) \log U / \chi^2_{2pq,\alpha}$ , and  $\chi^2_{2pq,\alpha}$  is the upper 100  $\alpha\%$  value of

$\chi^2$  with  $2pq$  degrees of freedom. The values of  $c_1$  are computed for  $\alpha = 0.005, 0.01, 0.025, 0.05, 0.1, 0.90, 0.95, 0.99, 0.995$ ,  $M = 1(1)10(2)20, 30, 60, 120$ , where  $M = n-p+1$ . These values are given in a technical report by Lee, Krishnaiah and Chang (1975). The upper 5% and 1% points are reproduced in Table 7 of this chapter. To check for the accuracy of the entries in Table 7, the above authors compared some of the values obtained by the Pearson type approximation with the corresponding exact values. These comparisons are given in Table 1.

Table 1  
Comparison of the Pearson type Approximation with exact expression for the Distribution of  $c_1$

p = 2 q = 3				p = 2 q = 20			
M	$\alpha$	L-K-C	Exact	$\alpha$	L-K-C	Exact	
1	0.05	1.286	1.289	0.05	1.928	1.932	
1	0.01	1.350	1.349	0.01	2.085	2.080	
5	0.05	1.029	1.029	0.05	1.243	1.243	
5	0.01	1.033	1.033	0.01	1.262	1.262	
9	0.05	1.010	1.011	0.05	1.129	1.128	
9	0.01	1.012	1.012	0.01	1.137	1.137	

The constant  $\alpha$  in Table 1 is defined by Eq. (3.3). Also, the values under the column "L-K-C" are the values of  $c_1$  obtained by Lee, Krishnaiah and Chang (1975) using the Pearson type approximation whereas the values under the column "Exact" are the corresponding values given by Gupta (1971). Table 1 indicates that the accuracy of the Pearson type approximation is sufficient for practical purposes.

#### 4. Test for Independence of Sets of Variates

Let  $\underline{z}' = (z'_1, \dots, z'_q)$  be distributed as a complex multivariate normal distribution with mean vector  $\mu' = (\mu'_1, \dots, \mu'_q)$  and covariance matrix  $\Sigma$ . Also, let  $E\{(z_i - \mu_i)(\overline{z_j - \mu_j})'\} = \Sigma_{ij}$ , and  $E(z_i) = \mu_i$ . It is assumed that  $z_i$  is of order  $p_i \times 1$  and  $p_1 + \dots + p_q = s$ . In this section we discuss the problem of testing the hypothesis  $H_2$  where

$$H_2: \Sigma_{ij} = 0 \quad (4.1)$$

for  $i \neq j = 1, \dots, q$ . Now let

$$A = \begin{bmatrix} A_{11} & A_{12} & \dots & A_{1q} \\ A_{21} & A_{22} & \dots & A_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ A_{q1} & A_{q2} & \dots & A_{qq} \end{bmatrix}$$

where

$$A_{gh} = \sum_{j=1}^N (z_{gj} - z_{g.})(\overline{z_{hj} - z_{h.}})', \quad z_{g.} = N^{-1} \sum_{j=1}^N z_{gj}$$

and  $(z'_{1j}, \dots, z'_{qj})$  is  $j$ -th independent observation on  $(z'_1, \dots, z'_q)$ .

The likelihood ratio statistic for testing  $H_2$  is

$$\lambda_2 = \frac{|A|}{\prod_{j=1}^q |A_{ij}|}. \quad (4.2)$$

In the analogous real case, Wilks (1935) derived the likelihood ratio test for multiple independence. The  $h$ th moment of  $\lambda_2$  is

$$E(\lambda_2^h) = \left[ \prod_{j=1}^s \frac{\Gamma(n+h-j+1)}{\Gamma(n-j+1)} \right] \left[ \prod_{i=1}^q \prod_{\alpha=1}^{p_i} \frac{\Gamma(n-\alpha+1)}{\Gamma(n+h-\alpha+1)} \right] \quad (4.3)$$

where  $n = N-1$  and  $\Gamma(\cdot)$  is the complete gamma function. The distribution of  $\lambda_2^{1/4}$  is approximated by Pearson's type I distribution with density

$$g(x) = \{\beta(\alpha+1, \epsilon+1)(\sigma_1 - \sigma_0)^{\alpha+\epsilon+1}\}^{-1} (x - \sigma_0)^\alpha (\sigma_1 - x)^\epsilon \quad (4.4)$$

where  $\sigma_0 < x < \sigma_1$  and  $\alpha$  and  $\epsilon$  are some real numbers.

Approximate percentage points of the distribution of  $\lambda_2 = -2 \log \lambda_2$  are constructed by Krishnaiah, Lee and Chang (1975, 1976) for  $p_i = p=1, 2, 3$ ;  $q=3, 4, 5$ ;  $\alpha = 0.01, 0.05, 0.10$ ;  $M = 1(1)20(2)30$  where  $M = n-s-3$ , and  $\Pr[\lambda_2 \leq c_2 | H_2] = (1-\alpha)$ . These percentage points are reproduced in Table 8. Percentage points for  $q=2$  are given in Table 7.

Now, consider a class of statistics  $W(0 \leq W \leq 1)$  whose moments are of the form

$$E\{W^h\} = K \frac{\prod_{j=1}^c y_j^a \prod_{k=1}^a x_k^a}{\prod_{j=1}^c \prod_{k=1}^a \Gamma[y_j(1+h) + y_j]} \frac{\prod_{k=1}^a \Gamma[x_k(1+h) + \xi_k]}{\prod_{j=1}^c \Gamma[y_j(1+h) + y_j]}, \quad h = 0, 1, \dots \quad (4.5)$$

where  $K$  is a normalizing constant such that  $E\{W^0\} = 1$  and  $\sum_{k=1}^a x_k = \sum_{j=1}^c y_j$ .

The likelihood ratio test statistics considered in this chapter are special cases of the above class of statistics. Box (1949) gave explicitly the first few terms of an asymptotic expression for the distribution of a class of statistics whose moments are of the form (4.5). But the first few terms alone are not sufficient to get the desired degree of accuracy in a number of practical situations. So, Lee, Krishnaiah and Chang (1976) gave terms up to order  $n^{-15}$  explicitly. In Table 2, given below, a comparison of the values obtained by using Pearson type approximation is made with the corresponding values obtained by the asymptotic expression of order  $n^{-13}$ .

Table 2

Comparison of the Pearson Type Approximation with the Asymptotic  
Expansion for the Distribution of  $\tilde{\lambda}_2$

n	q = 3 p = 1			q = 4 p = 2			q = 5 p = 1		
	c <sub>2</sub>	$\alpha_1$	$\alpha_2$	c <sub>2</sub>	$\alpha_1$	$\alpha_2$	c <sub>2</sub>	$\alpha_1$	$\alpha_2$
10	1.459	0.05	0.0499	-	-	-	4.011	0.05	0.0487
10	1.949	0.01	0.0100	-	-	-	4.811	0.01	0.0095
15	0.923	0.05	0.0500	5.733	0.05	0.0479	2.435	0.05	0.0497
15	1.233	0.01	0.0100	6.496	0.01	0.0093	2.914	0.01	0.0099
20	0.675	0.05	0.0500	3.958	0.05	0.0493	1.752	0.05	0.0499
20	0.902	0.01	0.0100	4.479	0.01	0.0098	2.096	0.01	0.0100
30	0.439	0.05	0.0501	2.455	0.05	0.0498	1.124	0.05	0.0500
30	0.587	0.01	0.0100	2.777	0.01	0.0099	1.344	0.01	0.0100

In this table,  $\alpha_1$  is the value of  $\alpha$  if we use the Pearson type approximation and  $\alpha_2$  is the value of  $\alpha$  if we use the asymptotic expression of order  $n^{-1/3}$ . From the table, we observe that the accuracy of the Pearson type approximation is sufficient for practical purposes.

## 5. Test for Sphericity

The likelihood ratio statistic for testing  $H_0: \Sigma = \sigma^2 \Sigma_0$  is given by

$$\lambda_3 = \frac{|A \Sigma_0^{-1}|}{(\text{tr } A \Sigma_0^{-1}/s)^s} \quad (5.1)$$

where  $\Sigma_0$  is known, A was defined in Section 4 and  $\text{tr } A$  denotes the trace of

A. The  $h^{\text{th}}$  moment of  $\lambda_3$  is known to be

$$E(\lambda_3^h) = \frac{s^{hs}}{\Gamma(sn + sh)} \prod_{j=1}^s \frac{\Gamma(n+h-j+1)}{\Gamma(n-j+1)} \quad (5.2)$$

Mauchly (1940) derived the likelihood ratio statistic for testing the hypothesis of sphericity when the underlying distribution is real multivariate normal.

The distribution of  $\lambda_3^{1/b}$  is approximated by a Pearson Type I distribution, where b is a suitably chosen integer. For  $M > 22$ , we took  $b=2$  and for  $M < 22$ ,  $b=4$ . Using the approximation described above, approximate upper percentage points of distribution of  $\tilde{\lambda}_3 = -2 \log \lambda_3$  were constructed by Krishnaiah, Lee and Chang (1975, 1976) for  $s = 2(1)10$ ,  $\alpha = 0.01, 0.05$ ,  $M = 1(1)20(2)30(5)50,60$ , where  $M = n-s-3$  and  $P[\tilde{\lambda}_3 \leq c_3 | H_0] = (1 - \alpha)$ . These values are reproduced in Table 9.

In Table 3, we compare the values obtained by the Pearson type approximation with the corresponding values obtained by using Box's asymptotic expression of order  $n^{-1/3}$ . Upper percentage points of the distribution of  $\tilde{\lambda}_3$  for  $\alpha = 0.01, 0.05$  and  $s = 3(1)6$  are also given by Nagarsenker and Das (1975).

Table 3

Comparison of the Pearson Type Approximation with the Asymptotic Expression  
for the Distribution of  $\tilde{\lambda}_3$

n	s = 5			s = 8		
	$c_3$	$\alpha_1$	$\alpha_2$	$c_3$	$\alpha_1$	$\alpha_2$
15	2.763	0.05	0.0496	-	-	-
15	3.265	0.01	0.0099	-	-	-
21	1.895	0.05	0.0498	4.565	0.05	0.0490
21	2.238	0.01	0.0100	5.093	0.01	0.0097
41	0.928	0.05	0.0500	2.160	0.05	0.0499
41	1.095	0.01	0.0100	2.409	0.01	0.0100
51	0.739	0.05	0.0500	1.711	0.05	0.0499
51	0.873	0.01	0.0100	1.908	0.01	0.0100

In the above table,  $\alpha_1$  is the value of  $\alpha$  obtained by using the Pearson type approximation whereas  $\alpha_2$  is the value of  $\alpha$  obtained by using Box's asymptotic expansion of order  $n^{-1/3}$ . This table indicates that the accuracy of Pearson type approximation is sufficient for practical purposes.

## 6. Test Specifying the Covariance Matrix

The modified likelihood ratio statistic for testing the hypothesis  $H_4: \Sigma = \Sigma_o$  is given by

$$\lambda_4 = (e/n)^{sn} |A\Sigma_o^{-1}|^n \text{etr}(-A\Sigma_o^{-1}). \quad (6.1)$$

The modified likelihood ratio test statistic is obtained from the likelihood ratio test statistic by changing N to n. The moments of  $\lambda_4$  are seen to be

$$E(\lambda_4^h) = (e/n)^{shn} |\Sigma_o|^{nh} |I + h\Sigma_o|^{-n(1+h)} \\ \times \prod_{i=1}^s \{\Gamma(n+nh+1-i)/\Gamma(n+1-i)\}. \quad (6.2)$$

Anderson (1958) derived the likelihood ratio statistic for testing the hypothesis that the covariance matrix is equal to a specified matrix when the underlying distribution is real multivariate normal. The distribution of  $\lambda_4^{1/b}$  can be approximated with a Pearson Type I distribution using the first four moments, where b is a suitably chosen integer. Using the above approximation, Krishnaiah, Lee and Chang (1975, 1976) computed the percentage points of the distribution of  $\tilde{\lambda}_4 = -2 \log \lambda_4$  for  $s = 2(1)10$ ,  $\alpha = 0.01, 0.05$ ,  $M=1(1)20(2)30$ , where  $M = n-s-1$  and  $P[\tilde{\lambda}_4 \leq c_4 | H_4] = (1-\alpha)$ . These percentage points are reproduced in Table 10.

## 7. Test for Multiple Homogeneity of the Covariance Matrices

Let  $\underline{z}_1, \dots, \underline{z}_q$  be independently distributed as complex  $p$ -variate normal with mean vectors  $\mu_1, \dots, \mu_q$  and covariance matrices  $\Sigma_{11}, \dots, \Sigma_{qq}$ , respectively. Also, let  $\underline{z}_{ij}$  ( $j = 1, \dots, N_i$ ) be the  $j^{\text{th}}$  independent observation on  $\underline{z}_i$ . In this section, we study the Pearson type approximation to the distribution of the likelihood ratio statistic for testing  $H_5$  where

$$H_5: \left\{ \begin{array}{l} \Sigma_{11} = \dots = \Sigma_{q_1 q_1} \\ \Sigma_{q_1 + 1, q_1 + 1} = \dots = \Sigma_{q_2^*, q_2^*} \\ \vdots \quad \vdots \quad \vdots \\ \Sigma_{q_{d-1}^* + 1, q_{d-1}^* + 1} = \dots = \Sigma_{qq} \end{array} \right.$$

$q_0^* = 0$ ,  $q_1^* = q_1$ ,  $q_d^* = q$  and  $q_j^* = q_1 + \dots + q_j$ . The modified likelihood ratio statistic (obtained by changing  $N_i$  to  $n_i$  in the likelihood ratio test statistic) for testing  $H_5$ , is given by

$$\lambda_5 = \frac{\prod_{i=1}^q |A_{ii}/n_i|^{n_i}}{\prod_{j=1}^d \left| \sum_{i=q_{j-1}^* + 1}^{q_j^*} A_{jj}/n_j^* \right|^{n_j^*}} \quad (7.1)$$

where  $n_i = N_i - 1$ ,  $n_j^* = \sum_{i=q_{j-1}^* + 1}^{q_j^*} n_i$  and

$$A_{ii} = \sum_{j=1}^{N_i} (z_{ij} - z_{i.})(\overline{z_{ij}} - \overline{z_{i.}})^T, \quad z_{i.} = \sum_{j=1}^{N_i} z_{ij}/N_i.$$

The moments of  $\lambda_5$  are given by

$$E(\lambda_5^h) = \left[ \frac{d}{\prod_{\alpha=1}^q n_\alpha} \right] \left[ \prod_{i=1}^p \prod_{\alpha=1}^{n_\alpha} \right] \frac{\Gamma(n_\alpha + h_n \alpha)}{\Gamma(n_\alpha + 1 - i)} \cdot$$

$$\times \frac{\Gamma(n_\alpha^{*+1-i})}{\Gamma(n_\alpha^{*+h_n \alpha^{*+1-i}})} . \quad (7.2)$$

Using the first four moments of  $\lambda_5$ , the distribution of  $\lambda_5^{1/b}$  can be approximated with a Pearson's Type I distribution where  $b$  is a suitably chosen integer. This approximation was used by Krishnaiah, Lee and Chang (1975, 1976) to compute approximate percentage points of the distribution of  $\tilde{\lambda}_5 = -2 \log \lambda_5$  for  $n_1 = n_0$ ,  $q = dk$  (i.e., there are  $k$  populations in each of the  $d$  groups). These points are reproduced in Table 11 for  $d=1$ .

In Tables 4 and 5, we compare the values obtained by the Pearson Type approximation for the distribution function of  $\lambda_5$  with the corresponding values obtained by using the Box's asymptotic expansion up to terms of order  $n^{-13}$ . In these tables, the constant  $c_5$  is defined as

$$P[-2 \log \lambda_5 \leq c_5 | H_5] = (1 - \alpha) . \quad (7.3)$$

Also,  $\alpha_1$  is the value of  $\alpha$  if we use the Pearson type approximation whereas  $\alpha_2$  is the value of  $\alpha$  if we use the asymptotic expression of order  $n^{-13}$ . Tables 4 and 5 indicate that the accuracy of the Pearson type approximation is sufficient for practical purposes.

In the real case, Wilks (1932) derived the likelihood ratio statistic for testing the homogeneity of the covariance matrices whereas Krishnaiah and Lee (1976) discussed how certain tests of hypotheses on linear structure of the covariance matrices can be reduced to the problem of testing for the multiple homogeneity of the covariance matrices.

Table 4

Comparison of the Pearson Type Approximation with the Asymptotic Expansion  
for the Distribution of  $\lambda_5$  when  $d = 1$

$n_0$	$q$	p = 3			p = 4		
		$c_5$	$\alpha_1$	$\alpha_2$	$c_5$	$\alpha_1$	$\alpha_2$
10	2	19.82	0.05	0.0501	33.00	0.05	0.0502
10	2	25.40	0.01	0.0100	40.21	0.01	0.0101
10	6	69.68	0.05	0.0500	12.14	0.05	0.0492
10	6	79.11	0.01	0.0100	13.39	0.01	0.0098
15	2	18.72	0.05	0.0501	30.33	0.05	0.0500
15	2	23.99	0.01	0.0100	36.93	0.01	0.0100
15	6	66.70	0.05	0.0500	113.77	0.05	0.0498
15	6	75.69	0.01	0.0100	125.48	0.01	0.0099
20	2	18.23	0.05	0.0500	29.18	0.05	0.0500
20	2	23.35	0.01	0.0100	35.52	0.01	0.0100
20	6	65.34	0.05	0.0499	110.45	0.05	0.0499
20	6	74.13	0.01	0.0100	121.78	0.01	0.0100

Table 5

Comparison of the Pearson Type Approximation with Asymptotic Expansion for the Distribution of  $\lambda_5$  when  $d > 1$

			p = 1			p = 2			p = 3			p = 4		
n <sub>0</sub>	q	d	c <sub>5</sub>	a <sub>1</sub>	a <sub>2</sub>	c <sub>5</sub>	a <sub>1</sub>	a <sub>2</sub>	c <sub>5</sub>	a <sub>1</sub>	a <sub>2</sub>	c <sub>5</sub>	a <sub>1</sub>	a <sub>2</sub>
10	6	3	8.01	0.05	0.0500	23.08	0.05	0.0500	46.97	0.05	0.0501	81.73	0.05	0.0503
10	6	3	11.62	0.01	0.0100	28.78	0.01	0.0100	55.03	0.01	0.0100	92.47	0.01	0.0101
10	6	2	9.70	0.05	0.0500	28.56	0.05	0.0500	58.66	0.05	0.0499	102.21	0.05	0.0500
10	6	2	13.57	0.01	0.0100	34.76	0.01	0.0100	67.44	0.01	0.0100	113.93	0.01	0.0100
20	6	3	7.91	0.05	0.0500	22.00	0.05	0.0500	43.22	0.05	0.0500	72.32	0.05	0.0500
20	6	3	11.49	0.01	0.0100	27.43	0.01	0.0100	50.60	0.01	0.0100	81.77	0.01	0.0100
20	6	2	9.59	0.05	0.0500	27.37	0.05	0.0500	54.49	0.05	0.0500	91.76	0.05	0.0500
20	6	2	13.42	0.01	0.0100	33.31	0.01	0.0100	62.63	0.01	0.0100	102.25	0.01	0.0100
30	6	3	7.88	0.05	0.0500	21.66	0.05	0.0500	42.12	0.05	0.0500	69.73	0.05	0.0500
30	6	3	11.44	0.01	0.0100	27.01	0.01	0.0100	49.31	0.01	0.0100	78.85	0.01	0.0100
30	6	2	9.56	0.05	0.0500	27.00	0.05	0.0500	53.26	0.05	0.0500	88.85	0.05	0.0500
30	6	2	13.38	0.01	0.0100	32.86	0.01	0.0100	61.22	0.01	0.0100	98.98	0.01	0.0100

## 8. Simultaneous Tests for the Homogeneity of Populations

In this section, we discuss the likelihood ratio test for the homogeneity of complex multivariate normal populations. The hypothesis of the homogeneity of the  $q$  complex multivariate distributions defined in Section 7 is equivalent to the hypothesis  $H_6$  where

$$H_6: \begin{cases} \Sigma_{11} = \dots = \Sigma_{qq} \\ u_1 = \dots = u_q. \end{cases} \quad (8.1)$$

The modified likelihood ratio statistic for testing  $H_6$  is given by

$$\lambda_6 = \frac{n^{pn} \prod_{i=1}^q |G_i|^{n_i}}{\prod_{i=1}^q n_i^{pn_i} |G + \sum_{i=1}^q N_i(z_{i.} - z_{..})'(z_{i.} - z_{..})|^n} \quad (8.2)$$

where  $n = \sum_{i=1}^q n_i$ ,  $n_i = N_i - 1$ ,

$$N = \sum_{i=1}^q N_i, z_{..} = \frac{1}{N} \sum_{i=1}^q \sum_{j=1}^{N_i} z_{ij}, z_{i.} = \frac{1}{N_i} \sum_{j=1}^{N_i} z_{ij}$$

$$G_i = \sum_{j=1}^{N_i} (z_{ij} - z_{i.})(\overline{z_{ij} - z_{i.}})', \quad G = \sum_{i=1}^q G_i.$$

The moments of  $\lambda_6$  are given by

$$E(\lambda_6^h) = [n^{phn} / \prod_{i=1}^q n_i^{phn_i}] \prod_{i=1}^p \prod_{j=1}^q \frac{\Gamma(n_j + h n_j + 1 - i)}{\Gamma(n_j + 1 - i)} \frac{\Gamma(n + q - i)}{\Gamma(n + h n + q - i)} \quad (8.3)$$

The distribution of  $\lambda_6^{1/b}$  is approximated with a Pearson Type I distribution where  $b$  is a suitably chosen integer. Using this approximation, percentage points of the distribution of  $\lambda_6 = -2 \log \lambda_6$  are computed by Chang, Krishnaiah and Lee (1975) for  $\alpha = 0.01, 0.025, 0.05, 0.10$ ;  $n_i = n_o; q = 2, 3, 4, 5$ ;  $p = 1, 2, 3, 4$  and  $M = n_o - p = 1(1) 20, 25, 30$ . Table 12 gives the value of  $c_6$  for  $\alpha = 0.05, 0.01$  where  $c_6$  is given by

$$P[\bar{\lambda}_6 \leq c_6 | H_6] = (1-\alpha). \quad (8.4)$$

To check for the accuracy of the entries in Table 12, we compared some of these values with the corresponding values obtained by using Box's asymptotic series of order  $n^{-1/3}$ . These comparisons are given in Table 6.

Table 6

Comparison of the Pearson Type Approximation with the Asymptotic Expansion

		p = 2			p = 3		
n <sub>0</sub>	q	c <sub>6</sub>	α <sub>1</sub>	α <sub>2</sub>	c <sub>6</sub>	α <sub>1</sub>	α <sub>2</sub>
7	3	28.03	0.05	0.0499	50.49	0.05	0.0497
7	3	34.15	0.01	0.0100	58.80	0.01	0.0098
10	3	27.45	0.05	0.0500	48.06	0.05	0.0498
10	3	33.42	0.01	0.0100	55.92	0.01	0.0099
15	3	27.04	0.05	0.0500	46.46	0.05	0.0499
15	3	32.90	0.01	0.0100	54.04	0.01	0.0100
20	3	26.84	0.05	0.0500	45.73	0.05	0.0500
20	3	32.67	0.01	0.0100	53.18	0.01	0.0100

In the above table,  $\alpha_1$  is the value of  $\alpha$  obtained by approximating  $\bar{\lambda}_6^{1/b}$  with Pearson's type I distribution whereas  $\alpha_2$  is the value of  $\alpha$  obtained by using Box's asymptotic series. Table 6 indicates that the accuracy of the values of  $\alpha_1$  is good.

## 9. Test Specifying the Values of the Covariance Matrix and Mean Vector

In this section, we consider the distribution of the likelihood ratio statistic for testing the hypothesis  $H_7$ , where

$$H_7: \begin{cases} \Sigma_{11} = \Sigma_o \\ \mu_1 = \mu_o \end{cases}$$

and  $\Sigma_o$  and  $\mu_o$  are known. The likelihood ratio statistic for testing  $H_7$  is given by

$$\lambda_7 = (e/N_1)^{pN_1} |G_1 \Sigma_o^{-1}|^{N_1} e^{\text{etr}[-\Sigma_o^{-1} \{G_1 + N_1(z_{11} - \mu_o)(\bar{z}_{11} - \mu_o)^T\}]} \quad (9.1)$$

The moments of  $\lambda_7$  are given by

$$E(\lambda_7^h) = \left(\frac{e}{N_1}\right)^{phN_1} \frac{1}{(1+h)^{pN_1(1+h)}} \prod_{i=1}^p \frac{\Gamma(N_1-i+N_1h)}{\Gamma(N_1-i)}. \quad (9.2)$$

Using the first four moments, Chang, Krishnaiah and Lee (1975, 1977) approximated the distribution of  $\lambda_7^{1/b}$  with Pearson Type I distribution where b is a suitably chosen integer. This approximation is used to compute the values of  $c_7$  for  $M=n_1-p-1 = 1(1)20(2)30$ ,  $n_1=N_1-1$  and  $p=2,3,4,5,6$  where

$$P[\tilde{\lambda}_7 \leq c_7 | H_7] = (1-\alpha).$$

and  $\tilde{\lambda}_7 = -2 \log \lambda_7$ . These values are given in Table 13 for  $\alpha = 0.05, 0.01$ .

## 10. Applications in Time Series in the Frequency Domain

In this section, we discuss as to how the likelihood ratio test procedures on the covariance matrices of the complex multivariate normal populations can be used in the area of inference on multiple time series.

Let  $\underline{X}'(t) = (X_1'(t), \dots, X_q'(t))$  ( $t = 1, \dots, T$ ) form a Gaussian, stationary, multiple time series with zero means and covariance matrix  $R(s) = (R_{jk}(s))$  where  $R_{jk}(s) = E\{\underline{X}_j(t) \underline{X}_k'(t + s)\}$  and  $\underline{X}_j(t)$  is of order  $p_j \times 1$ . The spectral density matrix of the above time series is given by  $F(\omega) = (F_{\ell j}(\omega))$  where

$$F_{\ell j}(\omega) = \frac{1}{2\pi} \sum_{s=-\infty}^{\infty} e^{-i\omega s} R_{\ell j}(s) . \quad (10.1)$$

A well known estimate (e.g. see Brillinger (1974) of  $F(\omega)$  is  $\hat{F}(\omega) = (\hat{F}_{\ell j}(\omega))$  where

$$\hat{F}_{\ell j}(\omega) = \frac{1}{(2m+1)} \sum_{r=-m}^{m} I_{\ell j}\left(\omega + \frac{2\pi r}{T}\right) \quad (10.2)$$

and  $m$  is a suitably chosen integer. In Eq. (10.2)

$$Z_{\ell}(\lambda) = \frac{1}{\sqrt{2\pi T}} \sum_{t=1}^T X_{\ell}(t) \exp(-it\lambda),$$

and

$$I_{\ell j}(\lambda) = Z_{\ell}(\lambda) \bar{Z}_{\ell}(\lambda).$$

Goodman (1963b) and Wahba (1968) showed that  $A(\omega) = (2m+1) F(\omega)$  is approximately distributed as complex Wishart distribution with  $(2m+1)$  degrees of freedom and  $E(\hat{F}(\omega)) = F(\omega)$ .

Now, let  $H_2(\omega)$ ,  $H_3(\omega)$  and  $H_4(\omega)$  denote the following hypotheses:

$$\begin{aligned}
 H_2(\omega): F_{\ell j}(\omega) &= 0 \quad (\ell \neq j = 1, \dots, q) \\
 H_3(\omega): F(\omega) &= \sigma^2 F_o(\omega) \\
 H_4(\omega): F(\omega) &= F_o(\omega)
 \end{aligned} \tag{10.3}$$

where  $\sigma^2$  is unknown and  $F_o(\omega)$  is known. Let the statistics  $L_2(\omega)$ ,  $L_3(\omega)$ , and  $L_4(\omega)$  be defined as follows:

$$L_2(\omega) = \frac{|A(\omega)|}{\prod_{i=1}^q |A_{ii}(\omega)|} \tag{10.4}$$

$$L_3(\omega) = \frac{|A(\omega) F_o^{-1}(\omega)|}{\{\text{tr } A(\omega) F_o^{-1}(\omega)/s\}^s} \tag{10.5}$$

$$L_4(\omega) = \left( e / (2m+1) \right)^{s(2m+1)} |A(\omega) F_o^{-1}(\omega)|^{(2m+1)} \tag{10.6}$$

$$\times \text{etr} (-A(\omega) F_o^{-1}(\omega)).$$

where  $(2m+1)A_{ij}(\omega) = \hat{F}_{ij}(\omega)$ . Also, let  $\tilde{L}_i(\omega) = -2 \log L_i(\omega)$  for  $i = 2, 3, 4$ .

The hypothesis  $H_2(\omega)$  is accepted or rejected according as

$$\tilde{L}_2(\omega) \leq d_2$$

where

$$P[\tilde{L}_2(\omega) \leq d_2 | H_2(\omega)] = (1 - \alpha). \tag{10.7}$$

We accept or reject  $H_3(\omega)$  according as

$$\tilde{L}_3(\omega) \leq d_3 \tag{10.8}$$

where

$$P[\tilde{L}_3(\omega) \leq d_3 | H_3(\omega)] = (1 - \alpha). \tag{10.9}$$

Similarly, the hypothesis  $H_4(\omega)$  is accepted or rejected according as

$$\tilde{L}_4(\omega) \leq d_4$$

where

$$P[\tilde{L}_4(\omega) \leq d_4 | H_4(\omega)] = (1 - \alpha). \quad (10.10)$$

Since  $A(\omega)$  is approximately distributed as the complex Wishart matrix, approximate values of  $d_2$ ,  $d_3$ , and  $d_4$  can be obtained from Table 8, Table 9, and Table 10, respectively.

$$\text{Next, let } H_2 = \bigcap_{j=1}^k H_2(\omega_j), H_3 = \bigcap_{j=1}^k H_3(\omega_j) \text{ and } H_4 = \bigcap_{j=1}^k H_4(\omega_j)$$

where  $\omega_1, \dots, \omega_k$  are widely separated. Then, we accept or reject  $H_2$  according as

$$T_2 \leq d_5 \quad (10.11)$$

where

$$P[T_2 \leq d_5 | H_2] = (1-\alpha) \quad (10.12)$$

and  $T_2 = \max(\tilde{L}_2(\omega_1), \dots, \tilde{L}_2(\omega_k))$ . An alternative procedure is to accept or reject  $H_2$  according as

$$T_2^* \leq d_6 \quad (10.13)$$

where

$$P[T_2^* \leq d_6 | H_2] = (1-\alpha), \quad (10.14)$$

and

$$T_2^* = \prod_{j=1}^k \tilde{L}_2(\omega_j).$$

Since  $\omega_1, \dots, \omega_k$  are widely separated,  $L_2(\omega_1), \dots, L_2(\omega_k)$  are distributed independently. So the critical values  $d_5$  and  $d_6$  can be computed by using the methods discussed in this chapter. We can propose similar procedures to test  $H_3$  and  $H_4$ .

Let  $H_5: F(\omega_1) = \dots = F(\omega_k)$ , where the frequencies  $\omega_1, \dots, \omega_k$  are "sufficiently" wide apart. Also, let  $\tilde{L}_5(\omega) = -2 \log L_5$ , where

$$L_5 = \frac{\prod_{i=1}^k |\hat{F}(\omega_i)|^{(2m+1)}}{\left| \sum_{i=1}^k \hat{F}(\omega_i)/k \right|^{k(2m+1)}}$$

Then, we accept or reject  $H_5$  accordingly as  $\tilde{L}_5 \leq d_5^*$  where

$$P[\tilde{L}_5 \leq d_5^* | H_5] = (1-\alpha).$$

Since  $\omega_1, \dots, \omega_k$  are "sufficiently" wide apart,  $\hat{F}(\omega_1), \dots, \hat{F}(\omega_k)$  are distributed independently. Also,  $(2m+1) \hat{F}(\omega_i)$  is distributed approximately as the complex Wishart matrix with  $(2m+1)$  degrees of freedom for  $i=1, \dots, k$ . Hence, the values of  $d_5^*$  can be obtained from Table 11.

Next, let  $\tilde{x}_i'(t) = (\tilde{x}_{i1}'(t), \dots, \tilde{x}_{iq}'(t))$ ,  $(t=1, \dots, T_i)$  be a Gaussian, stationary, multiple time series with zero means and covariance matrix  $R_i(s)$  and spectral density matrix  $F_i(\omega)$ , where  $R_i(s) = (R_{iuv}(s))$  and  $R_{iuv}(s) = E[\tilde{x}_{iu}(t) \tilde{x}_{iv}'(t+s)]$ .

Also, let  $x_1(t), \dots, x_k(t)$  be distributed independently and  $x_i(t)$  be of order  $p \times 1$  for  $i=1, \dots, k$ . Let the estimate  $\hat{F}_i(\omega)$  of  $F_i(\omega)$  be defined in a similar way as  $\hat{F}(\omega)$ . Here,  $(2m_i+1) \hat{F}_i(\omega)$  is distributed approximately as the complex Wishart matrix with  $2m_i+1$  degrees freedom. The hypothesis  $H_6(\omega): F_1(\omega) = \dots = F_k(\omega)$  is tested as follows. We accept or reject  $H_6$  accordingly as

$$\tilde{L}_6(\omega) \leq d_6^*$$

where

$$P[\tilde{L}_6(\omega) \leq d_6^* | H_6] = (1-\alpha),$$

$$\tilde{L}_6(\omega) = -2 \log L_6(\omega),$$

$$L_6(\omega) = \frac{\prod_{i=1}^k |\hat{F}_i(\omega)|^{(2m_i+1)}}{\left| \sum_{i=1}^k (2m_i+1)\hat{F}_i(\omega)/m_o \right|^{m_o}}$$

and  $m_o = 2(\sum_{i=1}^k m_i) + k$ . The critical values  $d_6^*$  can be obtained from Table 11 when the  $m_i$ 's are equal.

We will now illustrate the usefulness of some of the tables in this paper with vibration data on a C-5A transport aircraft.

Vibration measurements have been taken on the cargo deck of a C-5A transport aircraft to provide information about the dynamic environments that cargo must survive in transit and to understand better the distribution and transmission of vibrational energy throughout the aircraft structure. Measurements have been taken over certain periods by locating accelerometers at different locations on the cargo deck. We will treat each location as a variable. Data on the following variables were taken:

<u>Variables</u>	<u>Longitudinal Location</u>	<u>Lateral Location</u>	<u>Directional Orientation</u>
1 (FRV)	Forward	Right	Vertical
2 (FRL)	Forward	Right	Lateral
3 (FLV)	Forward	Left	Vertical
4 (FLL)	Forward	Left	Lateral
5 (ARV)	Aft	Right	Vertical
6 (ARL)	Aft	Right	Lateral
7 (ALV)	Aft	Left	Vertical
8 (ALL)	Aft	Left	Lateral

The basic unit of measurement is the acceleration due to gravity ( $g = 980 \text{ cm/sec}^2$ ). Let the spectral density of the data on the above 8 variables at frequency  $\omega$  be denoted by  $F(\omega)$  and let the corresponding population spectral density matrix be denoted by  $F(\omega)$ . The sample spectral density matrix at frequency  $\omega_j$  is  $F(\omega_j) = S_{j0} + iS_{j1}$  where  $\omega_1 = 0.15907\text{HZ}$ ,  $\omega_a = a\omega_1$ ,  $a = 2, 3, 4, 5$  and

$$S_{10} = \begin{bmatrix} .1112000 & .0027030 & .3204600 & .0033170 & .+22.000 & -.0007765 & .4451000 & -.0052160 \\ .0027030 & .0014000 & .0026990 & -.0004533 & .002720 & .0027110 & .0015183 & -.0031160 \\ .3204600 & .0016940 & .3316000 & .0036660 & .+340.000 & -.0006463 & .4600000 & -.0055720 \\ .0033170 & -.0004533 & .0036660 & .0004698 & .0047820 & -.0020620 & .6073768 & .0021640 \\ .+22.000 & .002720 & .+340.000 & .0047820 & .6171000 & -.0031030 & .6330000 & -.0031100 \\ .-0006463 & .0047820 & .-0020620 & .0051030 & .0080290 & -.0048130 & .-0030220 & .0008241 \\ .0015183 & .-0031030 & .6330000 & .0080290 & .-0048130 & .7303000 & .0008241 & .0106880 \\ .0016940 & .-0031100 & .-0030220 & .0008241 & .0106880 & .0008291 & .0106880 & .0008291 \end{bmatrix}$$

$$S_{11} = \begin{bmatrix} 0.0000000 & -.0000000 & -.0104100 & .0000439 & .0411000 & -.0152700 & .0657400 & .0159700 \\ .0000000 & 0.0000000 & .0000000 & .0000000 & .0000000 & -.0000000 & .0102700 & .0001463 \\ .0104100 & .0000000 & .0000000 & .0000000 & .1001000 & .-0172500 & .0005500 & .0181630 \\ .0000000 & .0000000 & .0000000 & .0000000 & .0011310 & .-000727 & .-0010270 & .0010250 \\ .-0000000 & .0000000 & .0000000 & .0000000 & .0000100 & .-0141800 & .0000000 & .0216900 \\ .0000000 & .0000000 & .0000000 & .0000000 & .0191600 & .0000000 & .0296300 & .-0.0111000 \\ .0102700 & .0000000 & .0000000 & .0000000 & .0063900 & .-0236300 & .0030000 & .0331130 \\ .0001463 & .0000000 & .0000000 & .0000000 & .-0211900 & .0011300 & .0111300 & .0000000 \end{bmatrix}$$

$$S_{20} = \begin{bmatrix} .0700803 & .0449563 & .0725760 & -.0001763 & .1112000 & .0004678 & .1664000 & -.00016200 \\ .0001609 & .0334357 & .0001718 & -.0003547 & -.0001157 & .043375 & -.002877 & -.0003396 \\ .0723740 & .0316719 & .0755300 & -.0003989 & .1121000 & .0036673 & .1056000 & -.00016970 \\ -.00013743 & -.0001357 & -.0001348 & .000466 & -.000482 & -.0003550 & -.0016440 & .0003926 \\ .1112000 & -.0001157 & .1123000 & -.00047842 & .2311000 & .0004914 & .2381000 & -.00024600 \\ .0001673 & .0013715 & .0003573 & -.0001550 & .0001916 & .0013363 & -.0014520 & .0001638 \\ .1064000 & .0003267 & .1090000 & -.0001090 & .2381000 & -.0015520 & .2754000 & -.00035660 \\ -.0001266 & .0003096 & -.00013976 & .0003192 & -.0024400 & .0001630 & .00305643 & .0017120 \end{bmatrix}$$

$$S_{21} = \begin{bmatrix} 0.000000 & 0.0010550 & -0.0024700 & -0.0020200 & 0.0015600 & 0.0023660 & 0.0065300 & -0.0027390 \\ -0.0010550 & 0.0000000 & -0.0011480 & 0.0000026 & -0.0023600 & 0.0001395 & -0.0020950 & -0.002051 \\ 0.0024700 & -0.0011480 & 0.0000000 & -0.0025000 & 0.0063300 & -0.0022350 & 0.052-000 & -0.0026400 \\ -0.0020200 & 0.0000026 & -0.0025000 & 0.0000000 & 0.0018100 & -0.0000834 & 0.0046600 & -0.0000812 \\ 0.0015600 & 0.0001395 & 0.0063300 & 0.0018100 & 0.0000000 & 0.00-24500 & 0.0130200 & -0.0050250 \\ 0.0023660 & -0.0022350 & -0.0000834 & -0.0024500 & 0.0000000 & 0.0000000 & -0.0029430 & -0.0000308 \\ 0.0065300 & 0.052-000 & 0.0046600 & 0.0000000 & 0.0000000 & 0.0000000 & 0.0000000 & -0.0038830 \\ -0.0027390 & -0.0020950 & -0.0000812 & 0.0000000 & 0.0000000 & 0.0000000 & 0.0000000 & 0.0000000 \end{bmatrix}$$

$$S_{30} = \begin{bmatrix} .0184644 & .0036411 & .0145700 & -.0001017 & .0153400 & .0011710 & .0120500 & -.0015350 \\ .0013811 & .0037033 & .0009133 & -.0000139 & .0037300 & -.0002562 & .0039800 & .0003206 \\ .0137743 & .0009133 & .0136600 & -.0000396 & .0171000 & .0012640 & .01225400 & -.0015900 \\ -.0005137 & -.03061394 & -.0306396 & .0003513 & -.0021510 & .0001656 & -.0037240 & -.0002224 \\ .0146440 & .0134300 & .0171000 & -.0029510 & .1321000 & .0018903 & .1422600 & -.0029800 \\ .0011713 & -.0102562 & .0012600 & .0001658 & .0014900 & .0015594 & .0018380 & .0017730 \\ .0022073 & .00139600 & .0224900 & -.0037200 & .1422600 & .0016380 & .1501000 & .0045980 \\ -.0013530 & .001393200 & .00015980 & -.0002224 & -.0042900 & -.00177300 & -.0C54500 & .0021300 \end{bmatrix}$$

$S_{31}$	$\begin{bmatrix} 0.0000000 & -0.1000000 & 0.0112290 & 0.000498 & -0.0022730 & 0.0014420 & 0.0030740 & -0.0016530 \\ 0.0000000 & 0.0000000 & 0.0002026 & 0.0000295 & 0.0021210 & 0.0006990 & -0.0011200 & -0.0007933 \\ -0.1112210 & -0.1112245 & 0.0000000 & 0.0001496 & 0.00021690 & 0.0011270 & 0.0077490 & -0.0015220 \\ -0.0018933 & -0.0330003 & -0.0001496 & 0.0000000 & -0.0020500 & -0.0006137 & -0.0019060 & 0.0007250 \\ -0.0022733 & -0.0021210 & -0.0021090 & 0.0020500 & 0.00001000 & 0.0005620 & 0.0006480 & -0.0005010 \\ -0.0014420 & -0.0030000 & 0.0001370 & 0.0000137 & -0.0060620 & 0.0000000 & -0.0075630 & 0.0031471 \\ -0.0000000 & -0.0011520 & -0.0017890 & 0.0011060 & 0.0020000 & 0.0075960 & 0.0010000 & -0.0005390 \\ -0.0011520 & -0.0017893 & 0.0001220 & -0.0000250 & 0.00083510 & -0.000171 & 0.0035390 & 0.0000000 \end{bmatrix}$
----------	---

<b>S<sub>40</sub></b>	[	.01751410	.00134323	.0173700	-.0004332	-.0014830	-.0020470	-.0015553	.0023876
		.00143249	.00144040	.00111940	-.00010330	.0034700	.00015400	.0027770	-.0031921
		.017176	.00111440	.00177400	-.00013300	.00614640	-.00164640	.00249440	.0016650
		.00143332	-.00141340	.00111360	.0011560	-.00034200	.00006001	-.0026070	-.0000072
		.00143640	.0036743	-.00134340	-.00034200	.00014900	.00217400	.0074600	.0010736
		.00204970	.0011542	.00016860	.00006001	.00274140	.00181600	.0031220	-.0020718
		.00335530	.0027770	.00030940	-.0026070	.0057400	.00312200	.0012000	-.0037500
		.0023873	-.00019461	.00146400	-.0000672	-.00034700	-.0020710	-.0037580	.0024530

$$S_{41} = \begin{bmatrix} 0.44444444 & 0.44444444 & -0.44444444 & -0.0003321 & 0.0241700 & 0.0001471 & 0.0239500 & 0.0000267 \\ -0.44444444 & 0.44444444 & 0.44444444 & 0.0002420 & 0.0171300 & 0.0010000 & 0.0017070 & -0.0011640 \\ 0.02233333 & 0.4444709 & 0.4444709 & -0.0027000 & 0.0251500 & 0.0000198 & 0.0257100 & 0.0001436 \\ 0.4444321 & -0.4442742 < 0 & 0.0002700 & 0.0000000 & -0.0021520 & -0.0011320 & -0.0429710 & 0.0136338 \\ -0.2323730 & -0.00117030 & -0.0251500 & 0.0023520 & 0.0000000 & 0.0062310 & 0.0078030 & -0.0005300 \\ -0.4444141 & -0.4441403 & 0.0001198 & 0.0011320 & -0.0062310 & 0.0000000 & -0.0056680 & -0.0000241 \\ -0.4444033 & -0.44410703 & -0.0257100 & 0.0024710 & -0.0073030 & 0.0055660 & 0.0000000 & -0.0001640 \\ -0.4444127 & 0.33116900 & 0.00011380 & -0.0010830 & 0.0085380 & 0.0000241 & 0.0081600 & 0.0000000 \end{bmatrix}$$

Let  $\hat{F}(\omega)$  be partitioned as

$$\hat{F}(\omega) = \begin{bmatrix} \hat{F}_{11}(\omega) & \hat{F}_{12}(\omega) & \hat{F}_{13}(\omega) & \hat{F}_{14}(\omega) \\ \hat{F}_{21}(\omega) & \hat{F}_{22}(\omega) & \hat{F}_{23}(\omega) & \hat{F}_{24}(\omega) \\ \hat{F}_{31}(\omega) & \hat{F}_{32}(\omega) & \hat{F}_{33}(\omega) & \hat{F}_{34}(\omega) \\ \hat{F}_{41}(\omega) & \hat{F}_{42}(\omega) & \hat{F}_{43}(\omega) & \hat{F}_{44}(\omega) \end{bmatrix}$$

where  $\hat{F}_{ii}(\omega)$  is of order  $2 \times 2$  for  $i = 1, 2, 3, 4$ . We computed  $\tilde{L}_2(\omega_j) = -2 \log L_2(\omega_j)$  where

$$L_2(\omega_j) = \frac{|(2m+1)\hat{F}(\omega_j)|}{\prod_{i=1}^4 |(2m+1)\hat{F}_{ii}(\omega_j)|},$$

and  $(2m+1) = 19$ . The values of  $\tilde{L}_2(\omega_j)$  in this case are found to be 47.760, 31.667, 33.684, 35.646 and 37.738 respectively. The value of the critical value  $d_2$  for  $n=19$ ,  $s=8$  and  $q=4$  from Table 8 is found to be 4.217 at 5% significance level. Since the computed values of  $L_2(\omega_j)$  are greater than the value from the table, we conclude that the sets (1,2), (3,4), (5,6), (7,8) of variables are not independent for each of the five frequencies considered.

Next, we computed the value of  $\tilde{L}_3(\omega_j) = -2 \log L_3(\omega_j)$

where

$$L_3(\omega_j) = \frac{|(2m+1)\hat{F}(\omega_j)|}{\{(2m+1)\text{tr } \hat{F}(\omega_j)/s\}^s}$$

$s=9$ ,  $(2m+1) = 19$ . It is found that the values of  $\tilde{L}_3(\omega_j)$ , in this case, are 78.321, 65.267, 63.226, 59.019 and 61.852 respectively. The critical value  $d_3$  for  $n=19$  and  $s=8$  is found to be 5.142 at 5% level. So, we

reject (individually) the hypotheses that  $F(\omega_j) = \sigma^2 I_p$  for  $j = 1, 2, 3, 4, 5$ .

For the hypothesis  $H_5$  we will consider only the first four variables.

Let spectral density matrix of the data on the first four variables at frequency  $\omega_1$  be denoted by  $\hat{F}(\omega_1)$  and let the corresponding population spectral density matrix be denoted by  $F(\omega_1)$ . The sample spectral density matrices at frequencies 0.15907 Hz, 0.47721 Hz and 0.79535 Hz are  $\hat{F}(0.15907) = S_{10} + i S_{11}$ ,  $\hat{F}(0.47721) = S_{20} + i S_{21}$  and  $\hat{F}(0.79535) = S_{30} + i S_{31}$ , respectively where

$$S_{10} = \begin{bmatrix} .3112000 & .0027030 & .3204000 & .0033170 \\ .0027030 & .0014000 & .0026990 & -.0008533 \\ .3204000 & .0026990 & .3316000 & .0036660 \\ .0033170 & -.0008533 & .0036660 & .0008898 \end{bmatrix}$$

$$S_{11} = \begin{bmatrix} 6.4200000 & -.0044000 & -.0104100 & .0006039 \\ .3664000 & 0.0000000 & .0049750 & .0003763 \\ .3104100 & -.0069750 & 0.0000000 & .0007700 \\ -.0004039 & -.0003763 & -.0007700 & 0.0000000 \end{bmatrix}$$

$$S_{20} = \begin{bmatrix} .0104600 & .0008811 & .0185700 & -.0005107 \\ .0008811 & .0007630 & .0009133 & .0006139 \\ .0185700 & .0009133 & .0196800 & -.0006396 \\ -.0005107 & -.0006139 & -.0006396 & .0005513 \end{bmatrix}$$

$$S_{21} = \begin{bmatrix} 0.0000000 & -.0006045 & .0012290 & .0004698 \\ .0006045 & 0.0000000 & .0002828 & .0000029 \\ -.0012290 & -.0002828 & 0.0000000 & .0001996 \\ -.0004698 & -.0000029 & -.0001996 & 0.0000000 \end{bmatrix}$$

$$S_{30} = \begin{bmatrix} .0177900 & -.0002297 & .0194000 & .0007346 \\ -.0002297 & .0006892 & -.0003287 & -.0006763 \\ .0194000 & -.0003287 & .0215900 & .0010720 \\ .0007346 & -.0006763 & .0010720 & .0007642 \end{bmatrix}$$

$$S_{31} = \begin{bmatrix} 0.0000000 & -.0004046 & .0002579 & .0004523 \\ .0004046 & 0.0000000 & .0003417 & -.0000041 \\ -.0002579 & -.0003417 & 0.0000000 & .0003401 \\ -.0004523 & .0000041 & -.0003401 & 0.0000000 \end{bmatrix}$$

We computed the value of  $L_5 = -2 \log L_5$  where

$$L_5 = \frac{\prod_{i=1}^3 |A(\omega_i)/(2m+1)|^{(2m+1)}}{|\sum_{i=1}^3 A(\omega_i)/3(2m+1)|^{3(2m+1)}},$$

$\omega_1 = 0.15907$  Hz,  $\omega_2 = 0.47721$  Hz,  $\omega_3 = 0.79535$  Hz,  $(2m+1) = 19$ . The value of  $L_5$  is found to be 224.72. The critical value  $d_5^*$  for  $n_0 = 19$ ,  $p = 4$  and  $k = 3$  from Table II is found to be 50.93 at the 5% significance level. Since  $L_5$  is greater than the value from the table, we conclude that the spectral density matrices  $F(\omega_1)$ ,  $F(\omega_2)$  and  $F(\omega_3)$  are significantly different from each other.

### References

- Anderson, T. W. (1958). An Introduction to Multivariate Statistical Analysis. Wiley, New York.
- Box, G. E. P. (1949). A general distribution theory for a class of likelihood criteria. Biometrika, 36, 317-346.
- Brillinger, D. R. (1974). Time Series: Data Analysis and Theory. Holt, Rinehart and Winston, Inc.
- Carmeli, M. (1974). Statistical theory of energy levels and random matrices in physics. J. Statist. Phy. 10, 259-297.
- Chang, T. C., Krishnaiah, P. R. and Lee, J. C. (1975). Approximations to the distributions of the likelihood ratio statistics for testing the hypotheses on covariance matrices and mean vectors simultaneously. Aerospace Research Laboratories TR 75-0176.
- Chang, T. C., Krishnaiah, P. R. and Lee, J. C. (1977). Approximations to the distributions of the likelihood ratio statistics for testing the hypotheses on covariance matrices and mean vectors simultaneously. In Applications of Statistics (P. R. Krishnaiah, ed.), pp. 97-108. North-Holland Publishing Company.
- Goodman, N. R. (1963a). Statistical analysis based on a certain multivariate complex Gaussian distribution (an introduction). Ann. Math. Statist. 34, 152-76.
- Goodman, N. R. (1963b). The distribution of the determinant of a complex Wishart distributed matrix. Ann. Math. Statist. 34, 178-180.
- Gupta, A. K. (1971). Distribution of Wilks' likelihood-ratio criterion in the complex case. Ann. Inst. Statist. Math. 23, 77-87.
- Hannan, E. J. (1970). Multiple Time Series. Wiley, New York.
- James, A. T. (1964). Distributions of matrix variates and latent roots derived from normal samples. Ann. Math. Statist. 35, 475-501.
- Khatri, C. G. (1965). Classical statistical analysis based on a certain multivariate complex Gaussian distribution. Ann. Math. Statist. 36, 98-114.
- Krishnaiah, P. R. (1976). Some recent developments on complex multivariate distributions. J. Multivariate Anal. 6, 1-30.
- Krishnaiah, P. R. and Lee, J. C. (1976). On covariance structure. Sankhyā, Series A, 38, 357-371.
- Krishnaiah, P. R., Lee, J. C. and Chang, T. C. (1975). On the distributions of the likelihood ratio statistics for tests of certain covariance structures of complex multivariate normal populations. Aerospace Research Laboratories TR 75-0169.

Krishnaiah, P. R., Lee, J. C. and Chang, T. C. (1976). The distributions of the likelihood ratio statistics for tests of certain covariance structures of complex multivariate normal populations. Biometrika 63, 543-549.

Lee, J. C., Krishnaiah, P. R. and Chang, T. C. (1975). Approximations to the distributions of the determinants of real and complex multivariate Beat matrices. Aerospace Research Laboratories TR 75-1068.

Lee, J. C., Krishnaiah, P. R. and Chang, T. C. (1976). On the distribution of the likelihood ratio test statistic for compound symmetry. S. African Statist. J. 10, 49-62.

Mauchly, J. W. (1940). Significance test for sphericity of a normal n-variate distribution. Ann. Math. Statist. 11, 204-209.

Nagarsenker, B. N. and Das, M. M. (1975). Exact distribution of sphericity criterion in the complex case and its percentage points. Comm. Statist. 4, 363-375.

Wahba, G. (1968). On the distributions of some statistics useful in the analysis of jointly stationary time series. Ann. Math. Statist. 39, 1849-1862.

Wilks, S. S. (1932). Certain generalization in the analysis of variance. Biometrika 24, 471-494.

Wilks, S. S. (1935). On the independence of K sets of normally distributed statistical variables. Econometrica 3, 309-325.

Wooding, R. A. (1956). The multivariate distribution of complex normal variables. Biometrika 43, 212-215.

## Appendix

Tables 7 through 13 give percentage points of various distributions discussed in this chapter. These tables are useful in implementation of the likelihood ratio test procedures. A description of these tables is given below.

Table 7: Percentage Points of the Distribution of the Determinant of the Complex Multivariate Beta Matrix

The entries in this table give the values of  $c_1$  where

$$P[C_1 \leq c_1] = (1 - \alpha),$$

$$c_1 = -\{(2n+q-p)\log U\}/\chi^2_{2pq,\alpha} \text{ and } U = |A_1(A_1+A_2)^{-1}|.$$

Here  $A_1$  and  $A_2$  are distributed independently as central  $p \times p$  complex Wishart matrices with  $n$  and  $q$  degrees of freedom respectively. These entries in this table are reproduced from a technical report by Lee, Krishnaiah and Chang (1975).

Table 8: Percentage Points of the Distribution of  $\tilde{\lambda}_2$  for Multiple Independence

The entries in this table give the values of  $c_2$  where

$$P[\tilde{\lambda}_2 \leq c_2] = (1 - \alpha),$$

$$\tilde{\lambda}_2 = -2 \log \lambda_2 \text{ and}$$

$$\lambda_2 = \frac{|A|}{\prod_{i=1}^q |A_{ii}|}$$

Here,  $A = (A_{ij})$  :  $pq \times pq$  is the central complex Wishart matrix with  $n$  degrees of freedom and  $E(A) = n\Sigma$ , and  $\Sigma = (\Sigma_{ij})$ . Also,  $A_{ij}$  and  $\Sigma_{ij}$  are of order  $p \times p$ . In addition,  $\Sigma_{ij} = 0$  for  $i \neq j = 1, 2, \dots, q$ ,  $M = n-s-3$  and  $s = pq$ .

Krishnaiah, Lee and Chang (1976) gave the values of  $c_2$  for  $\alpha = 0.05$ ,  $M = 1(1)4(2)16, 20, 24, 30$ ,  $p=1, 2, 3$ , and  $q=3, 4, 5$ . These entries are included in Table 8 with the kind permission of Biometrika Trustees. The remaining entries in the table are reproduced from a technical report by Krishnaiah, Lee and Chang (1975).

Table 9: Percentage Points of the Likelihood Ratio Test Statistic for Sphericity

The entries in this table are the values of  $c_3$  where

$$P[\tilde{\lambda}_3 \leq c_3] = (1 - \alpha),$$

$$\tilde{\lambda}_3 = -2 \log\{ |A\Sigma_o^{-1}| / (\text{tr} A\Sigma_o^{-1}/s)^s \},$$

and  $M=n-s-3$ . Also,  $A:s\times s$  is distributed as the central complex Wishart matrix with  $n$  degrees of freedom and  $E(A) = n\sigma^2\Sigma_o$  where  $\sigma^2$  is unknown and  $\Sigma_o$  is known. Upper 5% points of the distribution of  $\tilde{\lambda}_3$  are given in Krishnaiah, Lee and Chang (1976) for  $s=7(1)10$  and  $M=1(1)5, 7, 10, 15, 20, 30(5)50, 60$ . These entries are reproduced in Table 9 with the kind permission of Biometrika Trustees. The remaining entries in the table are reproduced from a technical report by Krishnaiah, Lee and Chang (1975).

Table 10: Percentage points of the Likelihood Ratio Statistic for Specifying the Covariance Matrix

The entries in this table give the values of  $c_4$  where

$$P[\tilde{\lambda}_4 \leq c_4] = (1 - \alpha),$$

$$\lambda_4 = (e/n)^{sn} |A\Sigma_o^{-1}|^n \text{etr}(-A\Sigma_o^{-1}) \text{ and } \tilde{\lambda}_4 = -2 \log \lambda_4.$$

In the above equation,  $A:s \times s$  is distributed as the central Wishart matrix with  $n$  degrees of freedom and  $E(A) = n\Sigma_0$  where  $\Sigma_0$  is known. Krishnaiah, Lee and Chang (1976) gave upper 5% points of the distribution of  $\tilde{\lambda}_4$  for  $s=2(1)7$ ,  $M=1(1)5, 7, 10, 15, 20, 30$  where  $M=n-s-1$ . These percentage points are reproduced in Table 10 with the kind permission of the Biometrika Trustees. The remaining entries in the table are reproduced from a technical report by Krishnaiah, Lee and Chang (1975).

Table 11. Percentage Points of the Likelihood Ratio Tests Statistic for the Homogeneity of the Covariance Matrices

The entries in this table are the values of  $c_5$  where

$$P[\tilde{\lambda}_5 \leq c_5] = (1 - \alpha),$$

$\tilde{\lambda}_5 = -2 \log \lambda_5$  and

$$\lambda_5 = \frac{\prod_{i=1}^q |A_{ii}/n_i|^{n_i}}{\left| \sum_{i=1}^q A_{ii}/n \right|^n}$$

where  $A_{11}, \dots, A_{qq}$  are distributed independently as central  $p \times p$  complex Wishart matrices with  $n_1, \dots, n_q$  degrees of freedom respectively,  $E(A_{11}/n_1) = \dots = E(A_{kk}/n_k)$  and  $n = n_1 + \dots + n_q$ . Upper 5% points of the distribution of  $\tilde{\lambda}_5$  are given in Krishnaiah, Lee and Chang (1976) for  $p = 3, 4$ ;  $q = 2(1)6, 8$ ;  $n_0 = 5(1)20, 25, 30$  where  $n_1 = \dots = n_q = n_0$ . These points are reproduced in Table 11 with the kind permission of Biometrika Trustees. The remaining entries in this table are reproduced from a technical report by Krishnaiah, Lee and Chang (1975).

Table 12: Percentage Points of the Likelihood Ratio Test Statistic  
for the Homogeneity of Complex Multivariate Normal Populations

The entries in this table are the values of  $c_6$  where

$$P[\tilde{\lambda}_6 \leq c_6 | H_6] = (1 - \alpha),$$

$\tilde{\lambda}_6 = -2 \log \lambda_6$  and  $\lambda_6$  is the likelihood ratio statistic for testing the hypothesis  $H_6$  of the homogeneity of  $q$  complex  $p$ -variate normal populations. The statistic  $\lambda_6$  is defined in Section 8. In the table,  $q$  denotes the number of populations  $M=N_0-p$  and  $N_0$  is the common size of various groups. Chang, Krishnaiah and Lee (1977) gave upper 5% points of the distribution of  $\tilde{\lambda}_6$  for  $q=2(1)5$  and  $M=1(1)20, 25, 30$ . These percentage points are reproduced in Table 12 with the kind permission of the North-Holland Publishing Company. The remaining entries in this table are reproduced from the technical report by Chang, Krishnaiah and Lee (1975).

Table 13: Percentage Points of the Likelihood Ratio Test for  $\Sigma = \Sigma_0$   
and  $\underline{\mu} = \underline{\mu}_0$ .

The entries in this table are the values of  $c_7$  where

$$P[\tilde{\lambda}_7 \leq c_7 | H_7] = (1 - \alpha), \quad \tilde{\lambda}_7 = -2 \log \lambda_7,$$

and  $\tilde{\lambda}_7$  is the likelihood ratio statistic for testing the hypothesis  $H_7$  where

$$H_7: \Sigma_1 = \Sigma_0, \quad \underline{\mu}_1 = \underline{\mu}_0.$$

Here  $\underline{\mu}_1$  and  $\Sigma_1$  are respectively the mean vector and covariance matrix of  $p$ -variate complex normal distribution and  $\underline{\mu}_0$  and  $\Sigma_0$  are known. Also,  $M=N_1-p-2$  and  $N_1$  is the sample size. Chang, Krishnaiah and Lee (1977) gave the values of  $c_7$  for  $\alpha = 0.05$ ,  $M = 1(1)20(2)30$  and  $p = 2(1)6$ . These values are reproduced in Table 13 with the kind permission of North-Holland Publishing Company. The remaining entries are reproduced from the technical report by Chang, Krishnaiah and Lee (1975).

Some of the entries in Tables 2-5 are reproduced from Krishnaiah, Lee and Chang (1976) with the kind permission of the Biometrika Trustees. Whereas some of the entries in Table 6 are reproduced from Chang, Krishnaiah and Lee (1977) with the kind permission of North-Holland Publishing Company.

Table 7  
Percentage Points of the Distribution of the Determinant of the Complex Multivariate Beta Matrix

M	$\alpha$	p = 3 q = 3		p = 3 q = 4		p = 3 q = 5		p = 3 q = 6		p = 3 q = 7		p = 3 q = 8	
		.050	.010	.050	.010	.050	.010	.050	.010	.050	.010	.050	.010
1	1.310	1.378	1.343	1.414	1.377	1.453	1.412	1.491	1.446	1.528	1.478	1.564	
2	1.132	1.153	1.154	1.177	1.177	1.202	1.201	1.228	1.225	1.253	1.249	1.278	
3	1.076	1.087	1.092	1.104	1.109	1.122	1.127	1.141	1.145	1.160	1.163	1.179	
4	1.050	1.057	1.062	1.070	1.075	1.083	1.089	1.098	1.103	1.113	1.117	1.128	
5	1.036	1.041	1.045	1.050	1.055	1.061	1.066	1.073	1.078	1.085	1.089	1.097	
6	1.027	1.030	1.034	1.038	1.043	1.047	1.051	1.056	1.061	1.066	1.071	1.077	
7	1.021	1.024	1.027	1.030	1.034	1.037	1.041	1.045	1.049	1.054	1.057	1.062	
8	1.017	1.019	1.022	1.024	1.028	1.030	1.034	1.037	1.041	1.044	1.048	1.052	
9	1.014	1.016	1.018	1.020	1.023	1.025	1.029	1.031	1.034	1.037	1.040	1.044	
10	1.011	1.013	1.015	1.017	1.019	1.021	1.024	1.026	1.029	1.032	1.035	1.037	
12	1.008	1.009	1.011	1.012	1.015	1.016	1.018	1.020	1.022	1.024	1.026	1.028	
14	1.007	1.007	1.009	1.010	1.011	1.012	1.014	1.015	1.017	1.019	1.021	1.022	
16	1.005	1.006	1.007	1.008	1.009	1.010	1.011	1.012	1.014	1.015	1.017	1.018	
18	1.004	1.005	1.006	1.006	1.007	1.008	1.009	1.010	1.011	1.012	1.014	1.015	
20	1.003	1.004	1.005	1.005	1.006	1.007	1.008	1.008	1.010	1.010	1.012	1.012	
30	1.002	1.002	1.002	1.002	1.003	1.003	1.004	1.004	1.005	1.005	1.006	1.006	
60	1.000	1.000	1.001	1.001	1.001	1.001	1.001	1.001	1.001	1.001	1.002	1.002	
120	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.001	1.001	
$\chi^2_{2pq,\alpha}$	28.8693	34.8053	36.4150	42.9798	43.7730	50.8922	50.9985	58.6192	58.1240	66.2062	65.1708	73.6826	

Table 7 (continued)

$M$	$\alpha$	$p = 3 \ q = 9$	$p = 3 \ q = 10$	$p = 4 \ q = 4$	$p = 4 \ q = 5$	$p = 4 \ q = 6$	$p = 4 \ q = 7$					
		.050	.010	.050	.010	.050	.010					
1	1.509	1.598	1.538	1.630	1.354	1.425	1.373	1.444	1.394	1.466	1.416	1.490
2	1.271	1.303	1.293	1.326	1.166	1.189	1.182	1.205	1.199	1.223	1.217	1.242
3	1.180	1.198	1.197	1.216	1.102	1.114	1.114	1.127	1.128	1.141	1.142	1.156
4	1.131	1.143	1.145	1.158	1.070	1.078	1.080	1.088	1.091	1.100	1.103	1.112
5	1.101	1.109	1.113	1.122	1.052	1.057	1.060	1.065	1.069	1.075	1.078	1.085
6	1.081	1.087	1.090	1.097	1.040	1.044	1.047	1.051	1.054	1.059	1.062	1.067
7	1.066	1.071	1.074	1.080	1.032	1.035	1.037	1.041	1.044	1.047	1.051	1.054
8	1.055	1.059	1.062	1.067	1.026	1.028	1.031	1.033	1.036	1.039	1.042	1.045
9	1.047	1.050	1.053	1.057	1.021	1.024	1.026	1.028	1.030	1.033	1.035	1.038
10	1.040	1.043	1.046	1.049	1.018	1.020	1.022	1.024	1.026	1.028	1.030	1.033
12	1.031	1.033	1.035	1.038	1.013	1.015	1.016	1.018	1.020	1.021	1.023	1.025
14	1.024	1.026	1.028	1.030	1.010	1.011	1.013	1.014	1.015	1.016	1.018	1.020
16	1.020	1.021	1.023	1.025	1.008	1.009	1.010	1.011	1.012	1.013	1.015	1.016
18	1.016	1.018	1.019	1.020	1.007	1.007	1.008	1.009	1.010	1.011	1.012	1.013
20	1.014	1.015	1.016	1.017	1.006	1.006	1.007	1.008	1.009	1.009	1.010	1.011
30	1.007	1.008	1.008	1.009	1.003	1.003	1.003	1.004	1.004	1.004	1.005	1.005
60	1.002	1.002	1.002	1.002	1.001	1.001	1.001	1.001	1.001	1.001	1.001	1.001
120	1.001	1.001	1.001	1.001	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
$\chi^2_{2pq,\alpha}$	72.1532	81.0688	79.0819	88.3794	46.1943	53.4858	55.7585	63.6907	65.1708	73.6826	74.4683	83.5134

Table 7 (continued)

M	$\alpha$	$P = 4, q = 8$		$P = 4, q = 9$		$P = 4, q = 10$		$P = 5, q = 5$		$P = 5, q = 6$		$P = 5, q = 7$	
		.050	.010	.050	.010	.050	.010	.050	.010	.050	.010	.050	.010
1	1.440	1.514	1.463	1.539	1.485	1.562	1.378	1.448	1.390	1.458	1.403	1.472	
2	1.235	1.261	1.254	1.280	1.272	1.299	1.191	1.214	1.203	1.226	1.216	1.240	
3	1.157	1.171	1.172	1.187	1.186	1.202	1.123	1.135	1.133	1.145	1.144	1.157	
4	1.115	1.124	1.127	1.137	1.139	1.149	1.087	1.095	1.096	1.104	1.105	1.114	
5	1.088	1.095	1.098	1.106	1.108	1.116	1.066	1.072	1.073	1.079	1.081	1.087	
6	1.070	1.076	1.079	1.084	1.088	1.094	1.052	1.056	1.058	1.062	1.065	1.069	
7	1.058	1.062	1.065	1.070	1.073	1.077	1.042	1.045	1.047	1.051	1.053	1.057	
8	1.048	1.052	1.055	1.058	1.061	1.065	1.035	1.037	1.039	1.042	1.044	1.047	
9	1.041	1.044	1.047	1.050	1.053	1.056	1.029	1.031	1.033	1.035	1.038	1.040	
10	1.035	1.038	1.040	1.043	1.046	1.048	1.025	1.027	1.028	1.031	1.032	1.034	
12	1.027	1.029	1.031	1.033	1.035	1.037	1.019	1.020	1.022	1.023	1.025	1.026	
14	1.021	1.023	1.025	1.026	1.028	1.030	1.015	1.016	1.017	1.018	1.020	1.021	
16	1.017	1.018	1.020	1.021	1.023	1.024	1.012	1.013	1.014	1.015	1.016	1.017	
18	1.014	1.015	1.017	1.018	1.019	1.020	1.010	1.011	1.011	1.012	1.013	1.014	
20	1.012	1.013	1.014	1.015	1.016	1.017	1.008	1.009	1.010	1.010	1.011	1.012	
30	1.006	1.006	1.007	1.008	1.008	1.009	1.004	1.004	1.005	1.005	1.006	1.006	
60	1.002	1.002	1.002	1.002	1.003	1.003	1.001	1.001	1.001	1.001	1.002	1.002	
120	1.001	1.001	1.001	1.001	1.001	1.001	1.000	1.000	1.000	1.000	1.000	1.000	
$\chi^2_{2pq,\alpha}$	83.6753	93.2168	92.8083	102.8160	101.8790	112.3290	67.5048	76.1539	79.0819	88.3794	90.5312	100.4250	

Table 7 (continued)

M	$\alpha$	p = 5 q = 8		p = 5 q = 9		p = 5 q = 10		p = 6 q = 6		p = 6 q = 7		p = 6 q = 8	
		.050	.010	.050	.010	.050	.010	.050	.010	.050	.010	.050	.010
1	1.419	1.488	1.436	1.505	1.453	1.522	1.392	1.459	1.400	1.465	1.418	1.474	
2	1.230	1.254	1.245	1.269	1.259	1.284	1.210	1.233	1.219	1.242	1.230	1.252	
3	1.156	1.169	1.168	1.182	1.180	1.194	1.140	1.152	1.148	1.161	1.158	1.171	
4	1.115	1.124	1.125	1.135	1.136	1.145	1.102	1.110	1.110	1.118	1.118	1.126	
5	1.089	1.096	1.098	1.105	1.107	1.114	1.079	1.084	1.085	1.091	1.092	1.098	
6	1.072	1.077	1.079	1.084	1.087	1.092	1.063	1.067	1.068	1.073	1.075	1.079	
7	1.059	1.063	1.066	1.070	1.072	1.077	1.051	1.055	1.056	1.060	1.062	1.066	
8	1.050	1.053	1.056	1.059	1.062	1.065	1.043	1.046	1.047	1.050	1.052	1.055	
9	1.042	1.045	1.048	1.050	1.053	1.056	1.037	1.039	1.040	1.043	1.045	1.047	
10	1.037	1.039	1.041	1.044	1.046	1.049	1.031	1.034	1.035	1.037	1.039	1.041	
12	1.028	1.030	1.032	1.034	1.036	1.038	1.024	1.026	1.027	1.029	1.030	1.032	
14	1.022	1.024	1.025	1.027	1.029	1.030	1.019	1.020	1.021	1.023	1.024	1.025	
16	1.018	1.019	1.021	1.022	1.024	1.025	1.015	1.017	1.018	1.018	1.020	1.021	
18	1.015	1.016	1.017	1.018	1.020	1.021	1.013	1.014	1.014	1.015	1.016	1.017	
20	1.013	1.013	1.015	1.015	1.017	1.017	1.011	1.011	1.012	1.013	1.014	1.014	
30	1.007	1.007	1.008	1.008	1.009	1.009	1.006	1.006	1.006	1.007	1.007	1.007	
60	1.002	1.002	1.002	1.002	1.003	1.003	1.002	1.002	1.002	1.002	1.002	1.002	
120	1.001	1.001	1.001	1.001	1.001	1.001	1.000	1.000	1.000	1.001	1.001	1.001	
$\chi^2_{2pq,\alpha}$		101.8790	112.3290	113.1450	124.1160	124.3420	135.8070	92.8083	102.8160	106.3950	117.8578	119.8710	131.1410

Table 7 (continued)

M	$\alpha$	P = 6 q = 9		P = 6 q = 10		P = 6 q = 7		P = 7 q = 8	
		.050	.010	.050	.010	.050	.010	.050	.010
1	1.421	1.485	1.433	1.497	1.401	1.464	1.406	1.467	
2	1.241	1.264	1.253	1.276	1.225	1.247	1.232	1.254	
3	1.168	1.180	1.178	1.191	1.154	1.167	1.162	1.174	
4	1.126	1.135	1.135	1.144	1.115	1.124	1.122	1.130	
5	1.100	1.106	1.107	1.114	1.090	1.096	1.096	1.102	
6	1.081	1.086	1.088	1.093	1.073	1.077	1.078	1.083	
7	1.068	1.071	1.074	1.078	1.060	1.064	1.065	1.069	
8	1.057	1.061	1.063	1.066	1.051	1.054	1.055	1.058	
9	1.049	1.052	1.054	1.057	1.044	1.046	1.047	1.050	
10	1.043	1.045	1.047	1.050	1.038	1.040	1.041	1.043	
12	1.033	1.035	1.037	1.039	1.029	1.031	1.032	1.034	
14	1.027	1.028	1.030	1.031	1.023	1.025	1.026	1.027	
16	1.022	1.023	1.025	1.026	1.019	1.020	1.021	1.022	
18	1.018	1.019	1.021	1.022	1.016	1.017	1.018	1.019	
20	1.016	1.016	1.018	1.018	1.014	1.014	1.015	1.015	
30	1.008	1.009	1.009	1.010	1.007	1.007	1.008	1.008	
60	1.002	1.003	1.003	1.003	1.002	1.002	1.002	1.003	
120	1.001	1.001	1.001	1.001	1.001	1.001	1.001	1.001	
$\chi^2_{2pq,\alpha}$	133.2570	145.0990	146.5670	158.9500	122.1080	133.4760	137.7010	149.7270	

TABLE 8  
Percentage Points of the Distribution  $\lambda_2$  for  
Multiple Independence

$N$	$\alpha$	P=1			P=2			P=3		
		10%	5%	1%	10%	5%	1%	10%	5%	1%
1	1.895	2.243	3.002	4.679	5.143	6.096	7.936	8.481	9.575	
2	1.607	1.902	2.543	4.065	4.488	5.314	7.061	7.542	8.584	
3	1.395	1.651	2.207	3.626	3.984	4.713	6.368	6.798	7.661	
4	1.233	1.459	1.949	3.264	3.584	4.230	5.802	6.193	6.974	
5	1.104	1.306	1.746	2.967	3.258	3.851	5.331	5.690	6.405	
6	1.001	1.184	1.581	2.722	2.987	3.530	4.933	5.265	5.925	
7	.914	1.081	1.445	2.513	2.759	3.259	4.592	4.899	5.512	
8	.842	.996	1.330	2.335	2.562	3.027	4.295	4.582	5.155	
9	.780	.923	1.233	2.180	2.393	2.826	4.035	4.305	4.841	
10	.727	.859	1.148	2.045	2.244	2.651	3.805	4.059	4.565	
11	.680	.805	1.075	1.926	2.113	2.495	3.600	3.840	4.310	
12	.639	.756	1.010	1.819	1.997	2.358	3.416	3.644	4.096	
13	.603	.713	.952	1.724	1.992	2.235	3.251	3.467	3.990	
14	.570	.675	.902	1.639	1.799	2.124	3.101	3.307	3.716	
15	.542	.641	.856	1.562	1.714	2.023	2.963	3.161	3.553	
16	.515	.610	.814	1.491	1.636	1.931	2.838	3.027	3.403	
17	.492	.582	.776	1.427	1.566	1.848	2.723	2.904	3.265	
18	.470	.556	.742	1.368	1.501	1.772	2.618	2.791	3.138	
19	.450	.532	.711	1.313	1.441	1.702	2.528	2.687	3.020	
20	.432	.511	.682	1.263	1.396	1.637	2.429	2.590	2.911	
22	.400	.473	.631	1.174	1.288	1.521	2.265	2.416	2.715	
24	.372	.439	.587	1.096	1.202	1.420	2.122	2.263	2.544	
26	.347	.410	.548	1.028	1.123	1.332	1.997	2.129	2.393	
28	.326	.386	.515	.968	1.062	1.254	1.866	2.010	2.259	
30	.307	.364	.485	.915	1.004	1.185	1.786	1.904	2.140	

TABLE 8 (Continued)

M	$\alpha$	P=1			P=2			P=3		
		10%	5%	1%	10%	5%	1%	10%	5%	1%
q=4										
1	2.962	3.364	4.234	7.374	7.909	9.963	12.428	13.049	14.276	
2	2.563	2.908	3.635	6.542	7.012	7.956	11.237	11.794	12.691	
3	2.248	2.551	3.186	5.886	6.306	7.149	10.271	10.775	11.768	
4	2.003	2.272	2.837	5.352	5.733	6.496	9.466	9.927	10.836	
5	1.887	2.049	2.556	4.909	5.259	5.955	8.783	9.210	10.049	
6	1.645	1.866	2.328	4.536	4.859	5.500	8.196	8.593	9.373	
7	1.511	1.713	2.138	4.217	4.514	5.111	7.604	8.057	8.785	
8	1.397	1.583	1.976	3.939	4.217	4.774	7.236	7.584	8.269	
9	1.298	1.472	1.837	3.697	3.958	4.479	6.837	7.166	7.812	
10	1.213	1.376	1.716	3.483	3.729	4.219	6.480	6.792	7.484	
11	1.139	1.291	1.610	3.293	3.525	3.986	6.161	6.457	7.037	
12	1.073	1.216	1.517	3.122	3.342	3.782	5.872	6.153	6.705	
13	1.014	1.149	1.434	2.969	3.178	3.596	5.608	5.878	6.404	
14	.961	1.090	1.359	2.830	3.029	3.427	5.369	5.625	6.129	
15	.914	1.036	1.292	2.704	2.894	3.273	5.149	5.395	5.878	
16	.871	1.007	1.231	2.588	2.770	3.133	4.946	5.182	5.646	
17	.831	0.943	1.176	2.482	2.656	3.005	4.758	4.986	5.632	
18	.796	0.902	1.125	2.384	2.552	2.886	4.585	4.805	5.234	
19	.763	0.865	1.079	2.294	2.455	2.777	4.424	4.636	5.050	
20	.733	0.831	1.036	2.211	2.366	2.676	4.274	4.479	4.878	
22	.680	0.770	0.960	2.061	2.205	2.493	4.004	4.194	4.568	
24	.633	0.717	0.694	1.930	2.064	2.335	3.765	3.944	4.297	
26	.593	0.672	0.838	1.814	1.941	2.195	3.553	3.723	4.054	
28	.557	0.631	0.767	1.712	1.832	2.071	3.364	3.525	3.839	
30	.525	0.595	0.742	1.621	1.734	1.961	3.195	3.346	3.645	

TABLE 8 (Continued)

N	$\alpha$	q=5						p=3					
		10%	5%	1%	10%	5%	1%	10%	5%	1%	10%	5%	1%
1	4.169	4.617	5.543	10.270	10.860	12.030	17.180	17.862	19.194				
2	3.623	4.011	4.011	9.222	9.746	10.784	15.717	16.330	17.534				
3	3.206	3.549	4.253	8.379	8.652	9.787	14.506	15.872	16.165				
4	2.677	3.163	3.813	7.683	8.115	8.968	13.483	14.002	15.012				
5	2.609	2.986	3.457	7.098	7.496	8.280	12.603	13.086	14.024				
6	2.388	2.642	3.163	6.599	6.967	7.94	11.637	12.269	13.166				
7	2.202	2.435	2.914	6.167	6.510	7.168	11.162	11.587	12.412				
8	2.042	2.259	2.703	5.789	6.110	6.745	10.563	10.965	11.743				
9	1.904	2.106	2.521	5.456	5.758	6.355	10.028	10.407	11.144				
10	1.764	1.973	2.361	5.159	5.444	6.009	9.547	9.906	10.606				
11	1.678	1.856	2.221	4.893	5.164	5.699	9.109	9.452	10.119				
12	1.585	1.752	2.096	4.654	4.912	5.420	8.713	9.041	9.677				
13	1.500	1.659	1.985	4.438	4.683	5.167	8.349	8.663	9.273				
14	1.425	1.575	1.885	4.241	4.475	4.938	8.015	8.316	8.901				
15	1.357	1.500	1.794	4.061	4.285	4.727	7.708	7.997	8.559				
16	1.295	1.431	1.713	3.895	4.111	4.534	7.423	7.702	8.242				
17	1.238	1.369	1.638	3.743	3.950	4.357	7.159	7.428	7.943				
18	1.186	1.312	1.569	3.602	3.801	4.192	6.914	7.173	7.676				
19	1.139	1.259	1.506	3.472	3.663	4.041	6.685	6.935	7.420				
20	1.095	1.210	1.448	3.350	3.536	3.900	6.471	6.713	7.183				
22	1.016	1.124	1.344	3.132	3.305	3.645	6.082	6.309	6.750				
24	.948	1.048	1.254	2.940	3.102	3.421	5.738	5.952	6.367				
26	.869	.983	1.176	2.771	2.924	3.224	5.430	5.633	6.026				
28	.836	.925	1.106	2.620	2.764	3.046	5.155	5.347	5.720				
30	.790	.873	1.045	2.485	2.622	2.891	4.906	5.089	5.444				

TABLE 9  
Percentage Points of the Likelihood Ratio Statistic for the  
Sphericity Test of Complex Covariance Matrix  
 $\alpha=0.05$

M	s=2	s=3	s=4	s=5	s=6	s=7	s=8	s=9	s=10
1	1.496	2.646	3.066	3.338	3.499	3.512	3.921	3.375	3.670
2	1.255	2.257	2.939	3.993	5.731	7.022	6.366	9.755	11.183
3	1.080	1.968	2.626	3.592	5.122	6.314	7.562	8.060	10.199
4	.949	1.745	2.374	3.265	4.633	5.740	6.906	8.122	9.383
5	.846	1.568	2.166	2.993	4.231	5.264	6.357	7.502	8.693
6	.763	1.423	2.006	2.993	3.895	4.863	5.891	6.973	8.100
7	.696	1.303	1.992	2.763	3.608	4.519	5.490	6.514	7.586
8	.639	1.201	1.844	2.567	3.362	4.222	5.142	6.115	7.137
9	.590	1.115	1.717	2.397	3.147	3.962	4.836	5.762	6.736
10	.543	1.040	1.606	2.248	2.959	3.732	4.565	5.440	6.301
11	.512	.975	1.509	2.116	2.791	3.528	4.322	5.168	6.061
12	.481	.916	1.423	1.999	2.642	3.346	4.104	4.915	5.773
13	.453	.866	1.346	1.895	2.508	3.181	3.908	4.606	5.511
14	.429	.820	1.277	1.801	2.387	3.031	3.730	4.476	5.272
15	.406	.779	1.215	1.715	2.277	2.836	3.567	4.287	5.053
16	.386	.741	1.158	1.638	2.177	2.772	3.419	4.113	4.851
17	.368	.707	1.107	1.567	2.086	2.659	3.281	3.952	4.666
18	.352	.676	1.060	1.503	2.001	2.554	3.155	3.803	4.494
19	.336	.648	1.016	1.443	1.924	2.457	3.038	3.665	4.335
20	.322	.622	.977	1.367	1.652	2.367	2.930	3.537	4.186
22	.298	.576	.906	1.269	1.724	2.207	2.735	3.306	3.918
24	.277	.536	.845	1.204	1.612	2.066	2.564	3.103	3.683
26	.258	.501	.791	1.129	1.513	1.943	2.413	2.925	3.474
28	.242	.471	.744	1.063	1.427	1.833	2.280	2.765	3.287
30	.228	.444	.702	1.005	1.349	1.735	2.160	2.622	3.120
35	.199	.388	.616	.883	1.188	1.531	1.909	2.322	2.769
40	.177	.345	.546	.787	1.061	1.370	1.711	2.084	2.488
45	.159	.311	.494	.710	.959	1.239	1.550	1.891	2.259
50	.144	.282	.450	.647	.875	1.132	1.417	1.730	2.069
60	.122	.239	.361	.549	.744	.964	1.209	1.478	1.771

TABLE 9 (Continued)

$\alpha = 0.01$								
M	s=2	s=3	s=4	s=5	s=6	s=7	s=8	s=9
1	2.174	3.440	4.746	6.112	7.531	8.995	10.501	12.041
2	1.623	2.930	4.095	5.326	6.620	7.967	9.361	10.794
3	1.570	2.554	3.602	4.724	5.912	7.158	8.455	9.797
4	1.378	2.263	3.218	4.247	5.346	6.505	7.716	8.976
5	1.229	2.033	2.908	3.659	4.880	5.962	7.101	8.286
6	1.109	1.845	2.653	3.537	4.491	5.506	6.579	7.708
7	1.010	1.689	2.439	3.265	4.160	5.117	6.129	7.192
8	927	1.957	2.258	3.032	3.875	4.779	5.739	6.750
9	957	1.445	2.102	2.831	3.627	4.484	5.397	6.360
10	797	1.348	1.966	2.655	3.409	4.223	5.093	6.013
11	745	1.263	1.847	2.499	3.216	3.992	4.623	5.703
12	699	1.188	1.741	2.361	3.044	3.785	4.579	5.423
13	658	1.121	1.647	2.238	2.889	3.598	4.360	5.170
14	622	1.062	1.563	2.126	2.750	3.429	4.160	4.940
15	590	1.009	1.486	2.026	2.626	3.276	3.979	4.738
16	560	960	1.418	1.934	2.508	3.135	3.813	4.537
17	534	917	1.355	1.850	2.402	3.007	3.660	4.359
18	510	877	1.297	1.774	2.305	2.868	3.520	4.195
19	488	840	1.244	1.703	2.216	2.779	3.369	4.043
20	468	806	1.195	1.638	2.133	2.677	3.267	3.901
22	432	746	1.108	1.522	1.985	2.495	3.050	3.646
24	402	694	1.034	1.421	1.856	2.337	2.860	3.423
26	375	650	968	1.333	1.743	2.197	2.691	3.225
28	352	610	910	1.255	1.643	2.072	2.542	3.049
30	331	575	859	1.106	1.554	1.962	2.409	2.892
35	289	503	753	1.042	1.368	1.731	2.129	2.561
40	256	447	671	929	1.222	1.549	1.908	2.298
45	230	402	604	838	1.104	1.401	1.729	2.085
50	209	366	550	764	1.007	1.280	1.560	1.907
60	177	309	466	649	.057	1.090	1.346	1.936

TABLE 10  
Percentage Points of the Likelihood Ratio Statistic for Specifying the  
Covariance Matrix

$\alpha = 0.05$

$H$	s=2	s=3	s=4	s=5	s=6	s=7	s=8	s=9	s=10
1	1.31	21.63	35.43	52.70	73.55	98.03	126.19	156.92	193.65
2	1.86	20.57	33.44	49.60	69.16	92.18	116.72	148.79	182.49
3	1.58	13.69	32.17	47.60	66.26	88.25	113.62	142.40	174.68
4	1.40	13.42	31.29	46.18	64.19	85.40	109.87	137.65	168.85
5	1.27	13.07	30.64	45.12	62.62	83.22	106.98	134.00	164.28
6	1.18	16.02	30.14	44.30	61.40	81.50	104.69	131.84	160.60
7	1.10	16.61	29.75	43.64	60.40	80.11	102.83	126.62	157.57
8	1.04	16.45	29.43	43.11	59.58	78.96	101.28	126.61	155.01
9	9.39	16.32	29.16	42.66	58.91	77.98	99.96	124.87	152.84
10	9.35	16.21	28.94	42.28	58.32	77.15	98.02	123.41	150.97
11	9.91	18.11	28.74	41.95	57.52	76.43	97.84	122.11	149.30
12	9.68	18.02	28.58	41.67	57.38	75.80	96.98	120.98	147.87
13	9.85	17.95	28.44	41.42	57.00	75.24	96.21	119.97	146.57
14	9.83	17.89	28.31	41.21	56.67	74.75	95.54	119.06	145.44
15	9.81	17.83	28.20	41.01	56.30	74.31	94.92	118.27	144.39
16	9.79	17.79	28.10	40.84	56.03	73.91	94.38	117.55	143.44
17	9.78	17.74	28.01	40.68	55.84	73.55	93.88	116.87	142.60
18	9.77	17.70	27.92	40.54	55.61	73.23	93.43	116.29	141.81
19	9.75	17.66	27.85	40.41	55.41	72.93	93.01	115.74	141.11
20	9.74	17.63	27.78	40.29	55.23	72.66	92.64	115.22	140.44
22	9.72	17.57	27.66	40.08	54.90	72.17	91.96	114.32	139.27
24	9.70	17.52	27.56	39.90	54.61	71.76	91.39	113.54	138.25
26	9.68	17.48	27.47	39.75	54.38	71.40	90.88	112.86	137.36
28	9.67	17.44	27.40	39.61	54.17	71.09	90.45	112.27	136.61
30	9.66	17.41	27.33	39.49	53.97	70.81	90.05	111.73	135.91

TABLE 10(Continued)

 $\alpha = 0.01$ 

M	S=2	S=3	S=4	S=5	S=6	S=7	S=8	S=9	S=10
1	15.90	27.97	43.50	62.62	95.36	111.79	141.94	175.84	213.42
2	15.23	26.46	40.91	58.73	90.00	104.77	133.08	165.01	200.54
3	14.84	25.53	39.23	56.25	76.51	100.11	127.13	157.63	191.59
4	14.58	24.91	36.18	54.51	74.02	96.77	122.81	152.21	184.99
5	14.39	24.47	37.37	53.23	72.15	94.24	119.52	148.06	179.91
6	14.25	24.13	36.74	52.24	70.71	92.25	116.89	144.72	175.77
7	14.14	23.86	36.25	51.44	69.54	90.63	114.77	142.02	172.38
8	14.05	23.65	35.85	50.80	58.59	89.30	112.99	139.70	169.53
9	13.98	23.47	35.52	50.26	67.79	88.18	111.48	137.78	167.15
10	13.92	23.33	35.24	49.80	67.11	87.22	110.20	136.15	165.05
11	13.86	23.20	35.00	49.41	66.51	86.39	109.09	134.71	163.21
12	13.83	23.10	34.80	49.08	56.01	85.67	108.12	133.44	161.62
13	13.80	23.00	34.63	48.76	55.56	85.03	107.25	132.28	160.20
14	13.76	22.92	34.47	48.52	55.17	94.48	106.50	131.28	158.91
15	13.73	22.85	34.33	48.29	54.82	83.96	105.82	130.62	157.76
16	13.71	22.79	34.20	48.08	54.49	83.52	105.20	129.58	156.76
17	13.68	22.72	34.09	47.89	54.21	83.11	104.64	128.86	155.81
18	13.67	22.67	33.99	47.73	53.95	82.74	104.13	128.18	154.94
19	13.65	22.62	33.90	47.57	63.71	82.39	103.66	127.58	154.14
20	13.63	22.58	33.81	47.43	53.50	82.09	103.24	126.99	153.41
22	13.60	22.51	33.67	47.18	63.12	81.53	102.48	126.00	152.15
24	13.58	22.44	33.54	46.97	62.73	81.06	101.83	125.13	151.03
26	13.55	22.39	33.43	46.79	62.51	80.65	101.27	124.37	150.03
28	13.54	22.34	33.34	46.63	52.27	80.30	100.77	123.72	149.20
30	13.52	22.30	33.26	46.49	52.05	79.98	100.33	123.14	148.44

TABLE 11  
 Percentage Points of Likelihood Ratio Statistic for the  
 Homogeneity of the Covariance Matrices

$\alpha=0.10$        $p=2$

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
4	10.07	16.50	22.99	28.93	34.72	40.41	46.01
5	9.49	15.94	21.89	27.60	33.17	38.63	44.02
6	9.15	15.43	21.23	26.80	32.22	37.56	42.82
7	8.92	15.08	20.78	26.26	31.00	36.84	42.01
8	8.76	14.84	20.47	25.87	31.15	36.33	41.43
9	8.63	14.65	20.23	25.58	30.31	35.93	41.00
10	8.54	14.51	20.04	25.36	30.54	35.63	40.66
11	8.46	14.40	19.89	25.18	30.33	35.39	40.39
12	8.40	14.30	19.77	25.03	30.16	35.19	40.17
13	8.35	14.23	19.67	24.91	30.02	35.03	39.98
14	8.30	14.16	19.59	24.80	29.89	34.89	39.82
15	8.27	14.10	19.51	24.71	29.79	34.77	39.69
16	8.23	14.05	19.45	24.64	29.70	34.67	39.57
17	8.21	14.01	19.39	24.57	29.62	34.57	39.47
18	8.18	13.97	19.34	24.51	29.54	34.49	39.38
19	8.16	13.94	19.30	24.46	29.48	34.42	39.29
20	8.14	13.91	19.26	24.41	29.43	34.36	39.23
25	8.06	13.79	19.11	24.23	29.22	34.12	38.96
30	8.02	13.72	19.02	24.11	29.08	33.96	38.77

$\alpha=0.10$        $p=3$

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
5	21.08	35.76	49.54	62.91	76.03	88.97	101.79
6	19.56	33.49	46.58	59.30	71.77	84.09	96.30
7	18.63	32.08	44.75	57.04	69.12	81.04	92.85
8	18.00	31.12	43.49	55.50	67.31	78.97	90.51
9	17.54	30.42	42.57	54.39	65.98	77.44	88.79
10	17.20	29.89	41.88	53.54	64.98	76.28	87.48
11	16.92	29.47	41.33	52.87	64.19	75.37	86.46
12	16.71	29.14	40.90	52.32	63.56	74.63	85.62
13	16.52	28.86	40.53	51.88	63.03	74.03	84.93
14	16.38	28.63	40.23	51.50	62.59	73.52	84.36
15	16.25	28.43	39.97	51.18	62.20	73.09	83.87
16	16.14	28.26	39.74	50.91	61.98	72.71	83.44
17	16.05	28.11	39.55	50.67	61.60	72.38	83.08
18	15.96	27.98	39.38	50.46	61.34	72.10	82.75
19	15.89	27.87	39.23	50.27	61.13	71.85	82.46
20	15.82	27.77	39.09	50.11	60.93	71.62	82.22
25	15.58	27.39	38.59	49.49	60.20	70.78	81.27
30	15.42	27.14	38.26	49.09	59.73	70.23	80.64

TABLE 11 (Continued)

 $\alpha=0.10 \quad p=4$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
6	36.34	62.39	87.08	111.15	134.35	158.32	181.59
7	33.48	58.07	81.43	104.21	126.56	148.87	170.92
8	31.68	55.33	77.82	99.78	121.41	142.82	164.09
9	30.43	53.42	75.31	96.68	117.74	138.61	159.27
10	29.52	52.01	73.45	94.39	115.04	135.46	155.73
11	28.82	50.93	72.02	92.62	112.34	133.05	153.00
12	28.27	50.07	70.59	91.23	111.28	131.13	150.80
13	27.82	49.37	69.96	90.08	109.32	129.54	149.05
14	27.45	48.80	69.19	89.14	108.78	128.24	147.57
15	27.15	48.31	68.54	88.33	107.95	127.16	146.32
16	26.88	47.89	67.99	87.65	107.04	126.23	145.26
17	26.66	47.53	67.51	87.06	106.33	125.42	144.35
18	26.45	47.22	67.09	86.54	105.71	124.69	143.54
19	26.28	46.94	66.73	86.09	105.17	124.08	142.83
20	26.13	46.70	66.41	85.68	104.70	123.53	142.20
25	25.55	45.80	65.21	84.19	102.92	121.47	139.90
30	25.19	45.22	64.43	83.24	101.79	120.16	138.41

 $\alpha=0.10 \quad p=5$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
7	56.07	97.04	136.08	174.28	211.34	249.31	286.40
8	51.49	90.09	126.96	163.03	198.64	233.96	269.00
9	48.53	85.58	121.00	155.68	189.91	223.87	257.52
10	46.47	82.38	116.75	150.43	183.59	216.64	249.45
11	44.93	79.98	113.59	146.52	179.02	211.24	243.26
12	43.75	78.14	111.12	143.46	175.38	207.09	238.56
13	42.79	76.66	109.14	141.02	172.47	203.72	234.69
14	42.03	75.44	107.53	139.00	170.11	200.93	231.59
15	41.39	74.43	106.18	137.33	168.11	198.61	228.94
16	40.85	73.57	105.04	135.91	166.41	196.68	226.71
17	40.39	72.83	104.06	134.69	164.96	194.99	224.83
18	39.98	72.20	103.20	133.62	163.69	193.53	223.18
19	39.64	71.64	102.46	132.70	162.58	192.22	221.69
20	39.33	71.14	101.79	131.97	161.61	191.08	220.42
25	38.20	69.35	99.39	128.89	158.03	186.95	215.70
30	37.49	68.21	97.87	126.98	155.77	184.34	212.71

TABLE 11 (Continued)

 $\alpha=0.05 \quad p=2$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
4	12.30	19.51	26.08	32.34	38.42	44.36	50.21
5	11.59	18.51	24.82	30.84	36.58	42.39	48.02
6	11.16	17.91	24.07	29.94	35.04	41.21	46.69
7	10.88	17.51	23.56	29.34	34.34	40.42	45.81
8	10.68	17.23	23.20	28.90	34.44	39.86	45.18
9	10.53	17.01	22.93	28.58	34.07	39.43	44.70
10	10.42	16.35	22.72	28.33	33.77	39.10	44.34
11	10.32	16.71	22.55	28.13	33.54	38.83	44.04
12	10.25	16.60	22.41	27.96	33.35	38.61	43.79
13	10.18	16.51	22.30	27.82	33.18	38.43	43.59
14	10.13	16.43	22.20	27.71	33.05	38.28	43.42
15	10.08	16.37	22.12	27.61	32.93	38.15	43.27
16	10.04	16.31	22.05	27.52	32.83	38.03	43.14
17	10.01	16.26	21.98	27.44	32.75	37.93	43.03
18	9.98	16.22	21.92	27.38	32.67	37.84	42.94
19	9.95	16.18	21.88	27.32	32.60	37.76	42.84
20	9.93	16.14	21.83	27.26	32.54	37.69	42.77
25	9.83	16.01	21.66	27.06	32.30	37.43	42.47
30	9.77	15.92	21.55	26.93	32.15	37.26	42.27

 $\alpha=0.05 \quad p=3$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
5	24.35	39.80	54.17	68.05	81.52	94.99	108.18
6	22.57	37.24	50.91	64.10	77.02	89.73	102.28
7	21.48	35.66	48.88	61.66	74.16	86.46	98.62
8	20.75	34.58	47.50	59.99	72.19	84.23	96.11
9	20.22	33.80	46.50	58.77	70.76	82.59	94.29
10	19.82	33.21	45.74	57.85	69.68	81.35	92.88
11	19.51	32.75	45.14	57.12	68.34	80.37	91.79
12	19.26	32.37	44.66	56.53	68.15	79.59	90.93
13	19.04	32.06	44.26	56.04	67.38	78.93	90.18
14	18.87	31.81	43.92	55.64	67.10	78.39	89.55
15	18.72	31.59	43.64	55.30	66.70	77.94	89.03
16	18.60	31.40	43.40	55.00	66.35	77.54	88.58
17	18.49	31.23	43.18	54.74	66.35	77.18	88.18
18	18.39	31.09	43.00	54.51	65.78	76.88	87.55
19	18.30	30.96	42.83	54.31	65.34	76.61	87.55
20	18.23	30.85	42.68	54.13	65.34	76.37	87.28
25	17.95	30.43	42.13	53.46	64.36	75.47	86.26
30	17.76	30.15	41.78	53.03	64.34	74.89	85.60

TABLE 11 (Continued)

 $\alpha=0.05$        $p=4$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
6	40.71	67.84	93.36	118.15	142.48	166.51	190.31
7	37.47	63.08	87.24	110.70	133.75	156.49	179.04
8	35.43	60.08	83.34	105.96	128.17	150.09	171.45
9	34.03	57.99	80.62	102.65	124.27	145.63	166.81
10	33.00	56.45	78.63	100.19	121.39	142.31	163.04
11	32.21	55.26	77.09	98.31	119.16	139.76	160.19
12	31.59	54.33	75.37	96.82	117.40	137.74	157.83
13	31.09	53.57	74.88	95.60	115.37	136.11	156.04
14	30.68	52.94	74.05	94.59	114.77	134.73	154.51
15	30.33	52.42	73.36	93.74	113.77	133.56	153.18
16	30.03	51.96	72.77	93.02	112.32	132.58	152.06
17	29.78	51.57	72.25	92.38	112.18	131.73	151.10
18	29.56	51.23	71.81	91.84	111.53	130.98	150.25
19	29.36	50.93	71.41	91.35	110.95	130.32	149.52
20	29.18	50.67	71.06	90.92	110.45	129.73	148.86
25	28.54	49.69	69.77	89.34	108.57	127.58	146.43
30	28.14	49.05	68.94	88.32	107.37	126.20	144.85

 $\alpha=0.05$        $p=5$ 

$n_o$	q=2	q=3	q=4	q=5	q=6	q=7	q=8
7	61.61	103.97	144.09	183.19	221.59	259.84	297.64
8	56.50	96.44	134.32	171.26	207.57	243.68	279.40
9	53.23	91.55	127.95	163.46	198.48	233.10	267.46
10	50.94	88.10	123.45	157.95	191.33	225.56	258.84
11	49.25	85.53	120.08	153.81	187.05	219.92	252.55
12	47.94	83.54	117.47	150.58	183.21	215.52	247.58
13	46.90	81.96	115.37	148.00	180.19	212.01	243.60
14	46.05	80.65	113.65	145.90	177.58	209.14	240.32
15	45.34	79.57	112.22	144.12	175.50	206.71	237.53
16	44.75	78.64	111.01	142.65	173.32	204.64	235.31
17	44.24	77.86	109.98	141.36	172.28	202.89	233.29
18	43.80	77.17	109.07	140.24	170.37	201.38	231.56
19	43.42	76.57	108.27	139.27	169.92	200.07	230.08
20	43.07	76.04	107.56	138.40	168.40	198.86	228.72
25	41.84	74.12	105.02	135.25	165.04	194.56	223.82
30	41.06	72.90	103.41	133.25	162.16	191.82	220.72

TABLE 12

Percentage Points of the Likelihood Ratio Test Statistic for the  
Homogeneity of Complex Multivariate Normal Populations

	$\alpha = 0.05$	$\beta = 1$		
	$q = 2$	$q = 3$	$q = 4$	$q = 5$
M				
1	8.12	12.47	16.36	20.06
2	8.03	12.49	16.50	20.30
3	7.97	12.51	16.59	20.45
4	7.94	12.52	16.64	20.55
5	7.92	12.53	16.68	20.62
6	7.91	12.53	16.71	20.67
7	7.89	12.54	16.73	20.71
8	7.89	12.55	16.75	20.74
9	7.88	12.55	16.77	20.77
10	7.87	12.56	16.78	20.79
11	7.87	12.56	16.79	20.81
12	7.86	12.56	16.80	20.82
13	7.86	12.56	16.81	20.84
14	7.86	12.56	16.81	20.85
15	7.85	12.57	16.82	20.86
16	7.85	12.56	16.83	20.87
17	7.85	12.57	16.83	20.88
18	7.85	12.57	16.84	20.89
19	7.85	12.57	16.84	20.89
20	7.85	12.57	16.85	20.90
25	7.84	12.57	16.86	20.92
30	7.84	12.58	16.87	20.94

TABLE 12 (Continued)

 $\alpha = 0.05$  $p = 2$ 

$M$	$q_1^* = 2$	$q_2^* = 3$	$q_3^* = 4$	$q_4^* = 5$
1	19.87	31.06	42.59	53.10
2	18.41	29.33	40.42	50.62
3	17.68	28.92	39.37	49.43
4	17.25	26.39	36.75	48.74
5	16.96	26.03	36.35	46.29
6	16.75	27.76	36.06	47.97
7	16.60	27.50	37.85	47.74
8	16.47	27.45	37.69	47.56
9	16.36	27.33	37.55	47.41
10	16.30	27.24	37.44	47.30
11	16.23	27.16	37.36	47.21
12	16.18	27.09	37.28	47.12
13	16.13	27.04	37.22	47.05
14	16.09	26.99	37.16	47.00
15	16.05	26.94	37.12	46.94
16	16.02	26.90	37.07	46.90
17	15.99	26.87	37.04	46.86
18	15.97	26.84	37.01	46.82
19	15.94	26.81	36.97	46.79
20	15.92	26.79	36.95	46.76
25	15.84	26.69	36.85	46.65
30	15.79	26.63	36.78	46.57

TABLE 12 (Continued)

 $\alpha = 0.05$  $p = 3$ 

$n$	$q_{\bar{B}} = 2$	$q_{\bar{B}} = 3$	$q_{\bar{B}} = 4$	$q_{\bar{B}} = 5$
1	30.46	59.39	80.89	101.75
2	32.90	54.56	74.91	94.65
3	31.04	52.04	71.30	90.97
4	29.90	50.49	69.88	88.70
5	29.13	49.42	68.37	87.16
6	28.56	48.05	67.62	86.05
7	28.13	48.06	66.90	85.21
8	27.79	47.61	66.34	84.55
9	27.52	47.24	65.89	84.02
10	27.30	46.94	65.52	83.57
11	27.11	46.66	65.19	83.21
12	26.96	46.40	64.94	82.90
13	26.82	46.27	64.71	82.63
14	26.70	46.11	64.50	82.40
15	26.60	45.97	64.33	82.20
16	26.51	45.84	64.17	82.01
17	26.42	45.73	64.04	81.85
18	26.35	45.63	63.92	81.71
19	26.28	45.54	63.80	81.58
20	26.22	45.45	63.70	81.46
25	25.99	45.14	63.31	81.01
30	25.83	44.92	63.05	80.70

TABLE 12(Continued)

 $\alpha = 0.05$  $p = 4$  $g_2 = 2 \quad g_3 = 3 \quad g_4 = 4 \quad g_5 = 5$ 

M	56.06	95.31	131.68	165.49
1	51.79	87.16	120.63	153.23
2	48.40	82.37	114.50	145.90
3	46.20	79.52	110.73	141.37
4	44.77	77.23	108.05	138.13
5	43.58	75.07	106.08	135.70
6	42.63	74.40	104.57	133.95
7	42.17	73.51	103.36	132.51
8	41.63	72.74	102.39	131.33
9	41.19	72.10	101.57	130.37
10	40.81	71.56	100.39	129.54
11	40.50	71.09	100.29	128.84
12	40.22	70.70	99.80	128.22
13	39.97	70.35	99.36	127.70
14	39.76	70.03	98.97	127.23
15	39.57	69.77	98.62	126.83
16	39.41	69.52	98.31	126.45
17	39.25	69.30	98.04	126.13
18	39.12	69.10	97.79	125.82
19	35.93	68.92	97.55	125.54
20	38.21	68.22	96.67	124.50
21	38.18	67.74	96.07	123.76

TABLE 12 (Continued)

 $\alpha = 0.01$  $p = 1$ 

M	$\theta = 2$	$\theta = 3$	$\theta = 4$	$\theta = 5$
1	11.78	16.66	20.99	25.06
2	11.65	16.59	21.15	25.34
3	11.57	16.71	21.25	25.52
4	11.53	16.72	21.32	25.63
5	11.50	16.73	21.37	25.72
6	11.48	16.74	21.40	25.78
7	11.45	16.75	21.43	25.83
8	11.45	16.75	21.46	25.87
9	11.44	16.76	21.47	25.90
10	11.43	16.76	21.49	25.92
11	11.42	16.77	21.51	25.95
12	11.42	16.77	21.52	25.96
13	11.41	16.77	21.52	25.98
14	11.41	16.77	21.53	26.00
15	11.40	16.78	21.54	26.01
16	11.40	16.78	21.55	26.02
17	11.40	16.78	21.55	26.03
18	11.40	16.78	21.56	26.04
19	11.39	16.78	21.57	26.05
20	11.39	16.78	21.57	26.06
25	11.38	16.79	21.59	26.09
30	11.38	16.79	21.60	26.11

TABLE 12 (Continued)

 $\alpha = 0.01$  $p = 2$ 

$M$	$q^2 = 2$	$q^2 = 3$	$q^2 = 4$	$q^2 = 5$
1	25.93	30.78	50.55	61.80
2	25.94	30.42	47.84	58.76
3	22.96	35.27	46.55	57.32
4	22.38	34.59	45.79	56.49
.5	21.99	34.15	45.30	55.96
6	21.72	33.83	44.95	55.58
7	21.51	33.60	44.70	55.30
8	21.35	33.42	44.49	55.08
9	21.23	33.27	44.34	54.92
10	21.12	33.16	44.21	54.78
11	21.04	33.05	44.10	54.66
12	20.97	32.98	44.02	54.57
13	20.90	32.90	43.94	54.49
14	20.84	32.84	43.87	54.42
15	20.80	32.79	43.82	54.36
16	20.76	32.74	43.77	54.31
17	20.72	32.70	43.72	54.26
18	20.69	32.67	43.68	54.22
19	20.66	32.63	43.64	54.18
20	20.63	32.60	43.62	54.14
25	20.53	32.48	43.49	54.02
30	20.45	32.40	43.41	53.93

TABLE 12 (Continued)

 $\alpha = 0.01$        $p = 3$ 

M	$\bar{g}_2 = 2$	$\bar{g}_3 = 3$	$\bar{g}_4 = 4$	$\bar{g}_5 = 5$
1	45.07	69.52	92.43	114.42
2	40.40	63.71	85.31	100.12
3	36.10	60.56	81.05	101.68
4	36.06	59.80	79.40	99.26
5	35.68	57.53	77.00	97.51
6	34.98	56.02	76.79	96.23
7	34.45	55.92	75.95	95.27
8	34.03	55.38	75.31	94.52
9	33.70	54.95	74.78	93.93
10	33.41	54.59	74.36	93.44
11	33.13	54.29	73.99	93.01
12	32.93	54.04	73.69	92.67
13	32.82	53.32	73.43	92.37
14	32.67	53.02	73.20	92.10
15	32.55	53.00	73.00	91.87
16	32.43	53.31	72.83	91.67
17	32.33	53.18	72.68	91.48
18	32.24	53.05	72.53	91.32
19	32.16	52.33	72.40	91.18
20	32.09	52.06	72.29	91.05
25	31.60	52.08	71.85	90.55
30	31.60	52.23	71.54	90.19

TABLE 12 (Continued)

 $\alpha = 0.01$  $p = 4$ 

$M$	$g_2 = 2$	$g_2 = 3$	$g_2 = 4$	$g_2 = 5$
1	64.34	109.42	146.98	183.36
2	61.52	99.01	134.17	168.24
3	57.35	93.43	127.28	160.05
4	54.75	84.99	122.89	154.88
5	52.95	87.44	119.37	151.30
6	51.53	85.95	117.35	148.66
7	50.62	84.27	115.94	146.65
8	49.62	83.18	114.59	145.05
9	49.13	82.29	113.48	143.74
10	45.64	81.56	112.57	142.68
11	45.21	80.34	111.80	141.76
12	47.82	80.41	111.16	140.99
13	47.49	79.95	110.58	140.31
14	47.20	79.56	110.09	139.73
15	46.95	79.20	109.66	139.21
16	46.72	78.90	109.28	138.77
17	46.53	78.62	108.94	138.36
18	46.34	78.36	108.62	138.00
19	46.18	78.15	108.35	137.66
20	46.03	77.94	108.09	137.37
21	45.80	77.74	107.80	136.20
30	45.07	76.60	106.42	135.40

TABLE 13  
Percentage Points of the Likelihood Ratio Test Statistic for  $\Sigma = \Sigma_0$  and  $\mu = \mu_0$

$H$	$\alpha = 0.05$						$\alpha = 0.01$					
	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$
1	29.33	34.13	51.51	72.40	91.05	110.61	26.52	42.19	61.37	86.17	110.61	110.61
2	19.20	32.16	48.40	68.30	91.05	103.56	24.99	39.56	57.46	78.79	103.56	103.56
3	19.51	30.91	46.33	65.10	87.12	98.92	24.05	37.94	54.96	75.27	98.92	98.92
4	18.03	30.02	44.97	63.02	84.25	95.57	23.41	36.02	53.24	72.79	95.57	95.57
5	17.93	29.37	43.91	61.72	82.09	92.36	22.36	36.00	51.95	70.93	93.03	93.03
6	17.43	28.87	43.09	60.22	80.36	92.36	22.61	35.37	50.95	69.47	91.82	91.82
7	17.22	28.47	42.43	59.23	78.96	92.36	22.34	34.67	50.15	68.38	89.41	89.41
8	17.05	28.15	41.83	58.41	77.80	92.12	21.47	34.47	49.38	67.35	88.97	88.97
9	16.32	27.69	41.44	57.73	76.84	91.94	21.94	34.14	48.96	66.94	86.96	86.96
10	15.91	27.55	41.07	57.15	76.00	91.79	21.65	33.85	48.58	65.85	85.99	85.99
11	15.71	27.46	40.73	56.64	75.29	91.65	21.65	33.62	48.11	65.27	85.17	85.17
12	15.53	27.23	40.45	56.20	74.64	91.55	21.41	33.41	47.77	64.76	84.44	84.44
13	16.55	27.15	40.20	55.82	74.09	91.46	21.23	33.23	47.46	64.31	83.61	83.61
14	15.43	27.02	39.93	55.48	73.61	91.37	21.17	33.07	47.22	63.92	83.26	83.26
15	16.44	26.91	39.79	55.17	73.15	91.30	21.10	32.93	46.98	63.56	82.74	82.74
16	16.38	25.91	39.61	54.90	72.76	91.23	21.03	32.81	46.77	63.25	82.29	82.29
17	15.34	26.72	39.45	54.65	72.40	91.17	21.07	32.69	46.56	62.96	81.81	81.81
18	16.30	26.63	39.31	54.43	72.06	91.12	21.02	32.59	46.41	62.70	81.51	81.51
19	16.26	25.50	39.19	54.22	71.76	91.07	21.07	32.50	46.26	62.46	81.16	81.16
20	16.23	26.49	39.00	54.04	71.51	91.03	21.03	32.41	46.11	62.24	80.86	80.86
22	15.17	26.37	38.85	53.71	71.02	90.95	20.95	32.26	45.87	61.86	80.31	80.31
24	16.12	26.27	38.77	53.43	70.60	90.98	20.98	32.14	45.65	61.53	79.83	79.83
26	16.07	26.16	38.52	53.18	70.25	90.83	20.83	32.03	45.47	61.25	79.42	79.42
28	16.04	26.10	38.13	52.93	69.93	90.76	20.76	31.94	45.31	61.08	79.07	79.07
30	16.01	25.03	38.27	52.79	69.65	90.65	20.74	31.85	45.17	60.79	78.74	78.74

TABLE 13 (Continued)

$\alpha = 0.025$						$\alpha = 0.10$						
$n$	1	2	3	4	5	6	1	2	3	4	5	6
11	23.05	24.75	26.44	28.13	29.82	31.51	37.74	39.43	41.12	42.81	44.50	46.19
12	18.90	18.80	18.70	18.60	18.50	18.40	30.02	30.97	31.92	32.87	33.82	34.77
13	18.65	18.65	18.65	18.65	18.65	18.65	29.72	29.72	29.72	29.72	29.72	29.72
14	18.50	18.50	18.50	18.50	18.50	18.50	29.60	29.60	29.60	29.60	29.60	29.60
15	18.53	18.53	18.53	18.53	18.53	18.53	29.48	29.48	29.48	29.48	29.48	29.48
16	18.49	18.49	18.49	18.49	18.49	18.49	29.38	29.38	29.38	29.38	29.38	29.38
17	18.43	18.43	18.43	18.43	18.43	18.43	29.29	29.29	29.29	29.29	29.29	29.29
18	18.39	18.39	18.39	18.39	18.39	18.39	29.21	29.21	29.21	29.21	29.21	29.21
19	18.35	18.35	18.35	18.35	18.35	18.35	29.13	29.13	29.13	29.13	29.13	29.13
20	18.28	18.28	18.28	18.28	18.28	18.28	29.00	29.00	29.00	29.00	29.00	29.00
21	18.23	18.23	18.23	18.23	18.23	18.23	28.99	28.99	28.99	28.99	28.99	28.99
22	18.18	18.18	18.18	18.18	18.18	18.18	28.79	28.79	28.79	28.79	28.79	28.79
23	18.14	18.14	18.14	18.14	18.14	18.14	28.71	28.71	28.71	28.71	28.71	28.71
24	18.10	18.10	18.10	18.10	18.10	18.10	28.63	28.63	28.63	28.63	28.63	28.63
$\Sigma$	125.4	125.4	125.4	125.4	125.4	125.4	30.7	30.7	30.7	30.7	30.7	30.7

END  
DATE  
FILMED

59 - 83

DTIC